

**SURVEYS OF THE FISH RESOURCES OF
THE WESTERN GULF OF GUINEA**

Guinea Bissau, Guinea, Sierra Leone and Liberia

Survey of the pelagic and demersal resources
29 April - 16 May 2006

Scientific Report DRAFT

Institute of Marine Research (IMR)
Norway

Ministry of Fisheries and Marine Resources
Sierra Leone

Centro de Investigação Pesqueira Aplicada (CIPA)
Guinea Bissau

Ministry of Fisheries and Marine Resources
Liberia

Centre National des Sciences Halieutiques (CNSHB)
Guinea

University of Ghana
Ghana

The DR FRIDTJOF NANSEN RESEARCH PROGRAMME is sponsored by the Norwegian Agency for Development Cooperation (NORAD). The Food and Agriculture Organization of the United Nations (FAO) provide support to the Programme through Project GCP/INT/730/NOR: International Cooperation with the Nansen Programme: Fisheries Management and Marine Environment. This project is the follow-up to the Project NORAD/FAO/UNDP GLO/92/013. The Institute of Marine Research (IMR), Bergen, Norway is responsible for the implementation of the Programme in cooperation with FAO Fisheries Department and the local fisheries administrations. The aim of the Nansen Programme is to assist developing countries in fisheries research, management and institutional strengthening.

The programme has previously focused on the mid to western Gulf of Guinea, but from 2004 surveys have covered the area, Part I from Côte d'Ivoire to Benin, and Part II from Nigeria to Gabon. The following surveys have been conducted in the Gulf of Guinea:

Area	Period
Cape Verga (Rep. of Guinea) to Cape St. Paul (Ghana)	02 - 25 June 1981
Togo to Cameroon	07 - 20 August 1981
Côte d'Ivoire and Ghana	12 - 20 October 1989
Benin, Togo, Ghana and Côte d'Ivoire	19 April - 06 May 1999
Benin, Togo, Ghana and Côte d'Ivoire	29 August - 17 September 2000
Benin, Togo, Ghana and Côte d'Ivoire	6 July - 09 August 2002
Benin, Togo, Ghana and Côte d'Ivoire (Gulf of Guinea Part I)	14 May - 08 June 2004
Nigeria, Cameroon, São Tomé and Principe (Gulf of Guinea Part II)	11 June – 13 July 2004
Benin, Togo, Ghana and Côte d'Ivoire (Gulf of Guinea Part I)	03 May - 29 May 2005
Nigeria, Cameroon, São Tom, Principe, Gabon and Congo	04 June – 15 July 2005

CRUISE REPORTS "DR. FRIDTJOF NANSEN"

**SURVEYS OF THE FISH RESOURCES OF
THE WESTERN GULF OF GUINEA
(Guinea Bissau, Guinea, Sierra Leone and Liberia)**

**Survey of the pelagic and demersal resources
29 April - 16 May 2006**

by

Ingvar Huse and Oddgeir Alvheim

Institute of Marine Research
P.O. Box 1870 Nordnes
N-5817 Bergen, Norway

Ibrahim Turay

Ministry of Fisheries and Marine Resources
Jomo Kenyatta Road, Brookfields,
Freetown, Sierra Leone

**Institute of Marine Research
Bergen, 2006**

TABLE OF CONTENTS

CHAPTER 1	INTRODUCTION.....	1
1.1	Objectives.....	1
1.2	Participation	1
1.3	Narrative.....	2
1.4	Survey effort.....	3
CHAPTER 2	METHODS.....	6
2.1	Meteorological and hydrographical sampling.....	6
2.2	Biological fish sampling.....	6
2.3	Zooplankton sampling.....	7
2.4	Benthos grab sampling	7
2.5	Biomass estimates	7
CHAPTER 3	OCEANOGRAPHIC CONDITIONS	11
3.1	Surface distribution	11
3.2	Vertical distribution.....	13
CHAPTER 4	RESULTS OF THE ACOUSTIC SURVEY.....	16
4.1	Guinea Bissau - Liberia.....	16
4.2	Review of results.....	22
CHAPTER 5	RESULTS FROM THE SWEPT AREA TRAWL SURVEY	23
5.1	Guinea Bissau.....	24
5.2	Guinea	28
5.3	Sierra Leone	31
5.4	Liberia	35
5.5	Swept area estimates	39
5.6	Overall review of results	40

Annex I	Records of fishing stations
Annex II	Length distributions of main species
Annex III	Summary of biological samples
Annex IV	Target species
Annex V	Swept-area biomass estimates
Annex VI	Estimates of sample variance
Annex VII	Instruments and fishing gear used
Annex VIII	Swept Area Estimates of Main Demersal Groups
Annex IX	Regional estimates

CHAPTER 1 INTRODUCTION

The survey of the Western part of Gulf reflects a discrepancy in coverage of the Western reaches of the Guinea Current LME, as this region was last surveyed in 1995. The present survey with R/V “Dr. Fridtjof Nansen” was initiated by the GCLME (Guinea Current Large Marine Ecosystem) and forms part of the cooperation between GCLME, FAO and IMR. The survey covered Guinea Bissau, Guinea, Sierra Leone and Liberia, and will continue along the continental shelf to include also the remaining GCLME countries.

The survey was organised by GCLME in cooperation with IMR and FAO under the FAO project GCP/INT/730/NOR: International cooperation with the Nansen Programme: Fisheries Management and Marine Environment, and the agreement between GCLME and IMR. This project is the continuation of a series of projects and agreements between NORAD, IMR and FAO involving surveys with the research vessel “Dr. Fridtjof Nansen”. The objectives of the survey was discussed and agreed upon during a pre-survey meeting held in Tema, Ghana, prior to the survey where representatives from GCLME and all countries surveyed were present together with representatives from FAO and IMR.

1.1 Objectives

Following the instructions from the GCLME and the recommendations from the pre survey meeting in Tema, Ghana, the main objectives of the survey were:

- to map the distribution and estimate the acoustic abundance of the main pelagic species / groups in the region
- to describe the distribution, composition and estimate the abundance of the main demersal species on the shelf by a swept-area trawl programme
- to collect bottom sediment samples to map the benthic biodiversity in the region.
- to collect phytoplankton and zooplankton samples for distribution and species identification
- to map the general hydrographic regime by using a CTD-sonde to monitor the temperature, salinity and oxygen at bottom trawl stations and on hydrographical transects
- on-the-job training on the main survey routines

1.2 Participation

Participants for the survey arrived in Conakry, Guinea 28 April. All participants stayed on board for the whole duration of the survey. The African participants represented the countries in the region covered by the survey, and one invited participant from Ghana.

From Guinea Bissau:

Luis Malabe da Fonseca, Vitorino Assau Nahada

From Guinea:

Amadou Bah (Team Leader), Amsoumane Keita, M'Mah Soumah

From Sierra Leone:

Sheku Sei (Team Leader), Ibrahim Turay (Local Cruise Leader)

From Liberia:

Alvin S. Jue-Seah, D.Wessey Kay

From Marine Fisheries Research Division, Tema, Ghana:

Emmanuel Lamptey

From Institute of Marine Research, Norway:

Oddgeir Alvheim, Ingvar Huse (Cruise Leader), Thor Egil Johansson and Tore Mørk.

1.3 Narrative

The vessel left Conakry (Guinea) at 17:30 on 28 April. The survey started at 21:00 the next day after the vessel arrived at the border between Senegal and Guinea Bissau at 17°09' W. The shelf was surveyed as much as possible during the day (0600 to 1800) by transects perpendicular to the general direction of the coastline, 30 or 40 NM (nautical miles) apart (30 NM transect distance was used in Guinea Bissau, but as this transect distance soon proved to exceed available time, 40 NM was adopted for the rest of the survey). Sampling was continued around the clock in the whole region, and for the whole duration of the survey.

The border to Guinea was reached on 05 May at 23:00h, and the coverage of Guinea was started immediately. The border between Guinea and Sierra Leone was reached on 07 May, and the vessel continued surveying into Sierra Leonean waters. The border between Sierra Leone and Liberia was reached 10 May, and the surveying continued through Liberian waters until the border with Cote d'Ivoire was reached 14 May at 17:00h, where the survey was discontinued. The vessel docked in Tema, Ghana on 16 May at 18:00h.

During the survey semi-random swept-area hauls were carried out on the shelf within the depth zones 20-30 m, 31-50 m, 51-100 m and when possible >100 m depth during daytime. Continuous acoustic recording and analysis was carried throughout the survey. Pelagic trawling on registrations and random blind hauls was carried out during dark hours when time permitted.

CTD-stations were taken at the bottom trawl stations. Zooplankton samples were taken either at the outer or at the inner end of each transect line with Hydrobios Multinet plankton sampler. At each plankton station a vertical net was also hauled from the bottom to the surface. Grab samples were taken at the innermost station of each transect. Additional grab samples were taken at every third transect line at the second innermost trawl station, and midway between the two innermost trawl stations.

1.4 Survey effort

Figure 1.1. shows the cruise tracks with bottom trawls, pelagic trawls and hydrographic stations, and Figure 1.2 shows the cruise tracks with plankton and grab stations.

Table 1.1 summarises the survey effort in each area.

Table 1.1 Number of hydrographic (CTD), Grab stations (G), plankton (P), pelagic trawl (PT) and bottom trawl (BT) stations, successful swept-area hauls, distance surveyed (NM) and size of survey area (NM²).




Figure 1.1 Course track with bottom trawl, pelagic trawl and hydrographic stations for Guinea Bissau-Guinea-Sierra Leone West (top) Sierra Leone East-Liberia (bottom). Depth contours are indicated.




Figure 1.2 Course track with plankton and grab sample stations for the survey area. Depth contours are indicated.

CHAPTER 2 METHODS

2.1 Meteorological and hydrographical sampling

Temperature, salinity and oxygen

CTD stations were taken in connection with all bottom trawl stations. Figure 1 presents positions for the CTD stations. A Seabird 911 CTD plus was used to obtain vertical profiles of temperature, salinity and oxygen. Real time plotting and logging was done using the Seabird Seasave software installed on a PC. The profiles were usually taken down to a few metres above the bottom, but normally not deeper than 300 m. The new oxygen sensor has shown to be very stable, and no calibration was conducted during the survey.

Thermosalinograph

The SBE 21 Seacat thermosalinograph was running routinely during the survey. Obtaining samples of sea surface salinity and relative temperature (5 m depth) every 10 sec during the survey.

Current speed and direction measurements (ADCP)

The ship-born Acoustic Doppler Current Profiler (ADCP) from RD Instruments was running throughout the survey. The ADCP was set to external trigger, triggered by the EK 500 system. The depth cell interval set to 8 m and the number of cells was set to 50.

Meteorological observations

Wind direction and speed, air temperature, global radiation and sea surface temperature (5 m depth) were logged automatically every nautical mile on an Aanderaa meteorological station.

2.2 Biological fish sampling

The trawl catches were sampled for species composition by weight and numbers. The deck sampling procedure is described in more detail by Strømme (1992). Length measurements (total length) were taken for most target species on most stations. The length of each fish was recorded to the nearest 1 cm below. The carapace length was measured to the nearest 0.5 cm below for shrimp. Basic information recorded at each fishing stations, i.e. trawl hauls, is presented in Annex I. Pooled length frequency distributions, raised to catch per hour, of selected species by area are shown in Annex II. Annex III gives basic information of all biological samples collected during the survey. Target groups for each country are indicated in Annex IV, while the swept-area estimates are presented in Annex V.

A description of the fishing gears used, acoustic instruments and their standard settings is given in Annex VII.

2.3 Zooplankton sampling

Zooplankton was collected with the Hydrobios Multinet zooplankton sampler that takes up to five discrete samples at predefined depths while measuring the water flow through the net. One sample was taken at one of the stations in each transect, either the outermost or the innermost, throughout the survey trying to cover both inshore and offshore areas. In addition a vertical net sample was taken for each Multinet station for comparison

2.4 Benthos grab sampling

The soft-bottom benthic macrofauna sampling was carried out using Peterson grab with a surface area of 0.20m². At each of the stations (Figure 1.2), the Peterson grab was deployed from an operated winch onto the seafloor. Five replicate samples were taken to obtain representative samples at each station, and to assess the patchiness in the distribution of the organisms. Two sediment replicates each were screened through sieves of mesh sizes 0.5 mm and 1.0 mm respectively to obtain adequate samples for both taxonomy and molecular analyses. The residue of the sieved sediment samples were fixed and put into plastic containers. Two of the sediment replicates were fixed in 10% borax pre-buffered formaldehyde while the other two were preserved in 90% ethanol. The ethanol in the samples were decanted and refilled with fresh ethanol solution after two days to avoid sample deterioration.

The containers were labelled according to the station numbers, replicate type, date, mesh size used, and the type of preservation used (e.g. N07A, 12/06/05, 0.5mm, Formaldehyde; C03D, 22/06/05, 1.0 mm, Ethanol). The samples were packed into boxes, for sorting and taxonomic identification on land. One replicate samples from all the stations were packed and sent to University of Ghana, Department of Oceanography & Fisheries while the three others were sent to Bergen Museum in Norway.

Additional sediment samples were taken at all the stations into ziplock bags, stored in a freezer and sent to Nigerian Institute for Oceanography and Marine Research, Lagos for both granulometric and chemical analyses.

Samples were taken at the innermost trawl station of each transect. Additional samples were taken for every third transect at the second innermost trawl station, and halfway between the innermost and the second innermost trawl station.

2.5 Biomass estimates

Acoustic abundance estimation

A SIMRAD EK500 Echo sounder was used and the echograms were stored on both paper and files. The acoustic biomass estimates were based on the integration technique. The Bergen

Integrator (BEI, see Knudsen 1990) was used for analysis and allocation of the integrated s_A -values (average area back scattering coefficient in m^2/NM^2) The splitting and allocation of the integrator outputs (s_A -values) was based on a combination of a visual scrutiny of the behaviour pattern as deduced from echo diagrams, the BEI analysis and the catch composition. The mean integrator value in each sampling unit (s_A -values) was divided between the standard categories/groups of fish, as noted below, on the basis of trawl catches and characteristics of echo traces:

- sardinella (*Sardinella aurita* and *S. maderensis*)
- PEL 1 (other clupeids than sardinella and anchovy)
- PEL 2 (carangids, scombrids, barracudas, hairtail)
- anchovy
- mesopelagic fish
- demersal fish
- plankton

The following target strength (TS) function was applied to convert s_A -values (mean integrator value for a given area) to number of fish (sardinella, anchovy, PEL 2):

$$TS = 20 \log L - 72 \text{ dB} \quad (1)$$

or in the form

$$C_F = 1.26 \cdot 10^6 \cdot L^{-2} \quad (2)$$

where L is total length and C_F is the reciprocal back scattering strength, or the so-called fish conversion factor. In order to split and convert the allocated s_A -values (m^2/NM^2) to fish densities (number per length group per NM^2) the following formula was used

where: N_i = number of fish in length group i

$$N_i = A \cdot s_A \cdot \frac{p_i}{\sum_{i=1}^n \frac{p_i}{C_{Fi}}} \quad (3)$$

A = area (NM^2) of fish concentration

s_A = mean integrator value (echo density) in area A (m^2/NM^2)

p_i = proportion of fish in length group i in samples from the area

C_{Fi} = fish conversion factor for length group i

The number per length group (N_i) was then summed and the total number of fish obtained:

$$N = \sum_{i=1}^n N_i \quad (4)$$

The length distribution of a given species within an area was computed by simple adding of the length frequencies obtained in the pelagic trawl samples within the area. In the case of co-occurrence of target species, the s_A value was split in accordance with length distribution and catch rate in numbers in the trawl catches. Biomass per length group (B_i) was estimated by applying measured weights by length (W_i) when available or theoretical weights (calculated by using condition factors), multiplied with number of fish in the same length group (N_i). The total biomass in each area was obtained by summing the biomass of each length group:

$$B = \sum_{i=1}^n N_i \bar{W}_i \quad (5)$$

The number and biomass per length group in each concentration were then added up to obtain totals for each region.

Biomass estimates based on swept-area method

In the bottom trawl survey, stock biomasses was estimated by the swept-area method with catch per haul as the index of abundance (see Strømme 1992). The general formula to estimate biomass B , using this method is:

$$B = \frac{A}{a} \cdot \frac{\bar{X}}{q} \quad (6)$$

A is the total area surveyed, a is the swept area of the net per haul, \bar{X} is the average catch per haul (the index of abundance) and q is the proportion of fish in the path of the net that are actually caught. The density of the resource is estimated as biomass per unit area. In a stratified survey of k non-overlapping strata, if the mean catch per haul in stratum i and its variance are denoted by \bar{X}_i and s_i^2 respectively, then an unbiased estimate of the population mean \bar{X} is the stratified mean \bar{X}_{st} , which is given by:

$$\bar{X}_{st} = \frac{1}{N} \sum_{i=1}^k N_i \bar{X}_i = \sum_{i=1}^k W_i \bar{X}_i \quad (7)$$

where $W_i = \frac{N_i}{N} = \frac{A_i}{A}$ is the relative size of the i^{th} stratum (A_i is the area of the i^{th} stratum and

A is the total area surveyed). The variance of the stratified mean is given by

$$\text{var}(\bar{X}_{st}) = \sum_{i=1}^k W_i^2 \text{var}(\bar{X}_i) = \sum_{i=1}^k W_i^2 \frac{s_i^2}{n_i} \quad (8)$$

where n_i is number of hauls in the i^{th} stratum and n is the total number of hauls in the survey.

Table 1.1 shows the areas used in the swept-area method to estimate biomass for the different regions. A stratified semi-random design was used with depth and country as stratification factors. Estimated total biomass by species/group was obtained by summing estimates for each depth stratum.

For conversion of catch rates (kg/hour) to fish densities (t/NM^2), the effective fishing area was considered as the product of the wing spread and the haul length, or distance over the bottom, as measured by means of the SCANMAR® equipment based on GPS readings. The area swept for each haul was thus 18.5 m (traditionally applied wing spread for the “Nansen” bottom trawl) times the distance trawled, raised to $NM^2/hour$. The catchability coefficient (q), i.e the fraction of the fish encountered by the 18.5 m horizontal opening of the trawl that was actually caught, was assumed equal to 1 for comparison with previous surveys. Mean fish densities by species and strata, were calculated by the swept-area module in NAN-SIS.

Total biomass estimates by species and their confidence intervals were obtained from a stratified mean density estimator (using equations 1, 2, and 4 in ANNEX VI on a spreadsheet, ANNEX VII) and raised to total area. Since NAN-SIS does not produce variance estimates of the mean densities (ANNEX IV), the 95% confidence limits for the biomass estimates were calculated with the underlying assumption that the coefficient of variation ($CV = SD/mean$) is constant when catch rates in kg/hour are converted to densities (t/NM^2). In other words the area swept (normalised per hour) was approximately constant for each haul. Coefficients of variation of the catch rates, by depth strata for each species or group, were obtained using the WinGrafer module of NAN-SIS. Variance of the densities were estimated from the mean and the CV , and equations 2, 3, 6 and 7 in ANNEX V were used to calculate standard error (SE) on the arithmetic mean and confidence intervals (see the spreadsheet BIOMASS.xls, and example in ANNEX VII). GRAFER was also used to produce the figures and tables with grouped catch-rates and time-series presented in this report. SE and confidence intervals in the figures are based on the arithmetic mean, but the lognormal based Pennington’s estimator can also be calculated (equations 8 to 12 in ANNEX VI).

CHAPTER 3 OCEANOGRAPHIC CONDITIONS

3.1 Surface distribution

The surface layer temperature (5 m depth) was continuously recorded during the cruise. Figure 3.1 shows the horizontal distribution of sea surface temperature (SST) for the survey area.

Guinea Bissau and Guinea

The sea surface temperatures outside of Guinea Bissau and Guinea were characterised by water masses with temperatures ranging from 22°C in the Northwest to 29°C in the Southeast towards the border of Sierra Leone. This water was assumed to be influenced by water from the Canary Current as an upwelling flowing in across the wide shelf area in this region. A local participating scientists from Guinea (Dr. Amsoumane Keita, Pers. Comm.) who had taken a PhD on this phenomenon had found that this upwelling was related to the rainy season.

On the Southeastern part of the wide shelf area towards Sierra Leone, tropical water masses (>29°C) were encountered.

Sierra Leone and Liberia

Here the tropical water masses (>29°C) dominated the SST throughout the area.




Figure 3.1 Horizontal distribution of surface temperature (5 m depth) in the survey area.

The surface salinity (Figure 3.2 a and b) was recorded from the Thermosalinograph at 5 m depth. The salinity was characterised by oceanic water in the West, and by coastal influence in the East.

Guinea Bissau and Guinea

Generally the salinity was corresponding with temperature in these countries, in the sense that the upwelling of deeper oceanic water was also identifiable in the salinity. In Guinea Bissau the upwelling could be traced directly by a deep intrusion at 35.9 ppt, being reduced to 35.5 ppt towards the shore. These saline water masses also penetrated from the deep into most of the Guinean shelf, with salinities of 35.5 ppt still measured at the Western shelf of Sierra Leone.

Sierra Leone and Liberia

The sea surface salinity in these two countries was more typical of a coastal narrow shelf tropical situation, with salinities ranging between 34.7 and 35.3 ppt, decreasing even further to 34.1 ppt in areas with river outlets (e. g. Southern Liberia).





Figure 3.2 Horizontal distribution of surface salinity (5 m depth) at a) Guinea Bissau - Guinea b) Sierra Leone – Liberia.

3.2 Vertical distribution

Figures 2.3a-d shows the vertical distribution of temperature, salinity and dissolved oxygen as recorded on the hydrographic transects worked during the survey.



a) Guinea Bissau




Figure 3.3a-d Vertical sections of temperature, salinity and oxygen.

The temperature and salinity data indicate that the Western upwelling does not originate from very deep water, rather from intermediate depths (<200 m), as the deeper water masses are less saline than the upper, and are only balanced density wise by the positive temperature gradient towards the surface.

In the South, the surface water is influenced by fresh water from the coast, and the deepest water is the most saline in theis area.

The oxygen data indicate that most of the oxygen content in the water originates from surface diffusion, creating favourable oxygen conditions e. g. on the Guinean shelf. Deeper water masses are, however, poor in oxygen, probably due to high degradation of biological material.

Wind conditions

Figure 3.3 shows wind conditions during the survey. There was little wind and prevailing good weather all through the survey period. Whatever wind we had was northerly in the Northwest, a gust of easterly wind in the middle and a south-westerly gust at the end of the survey. But generally we had calm winds and very favourable sea conditions all through the survey.




Figure 3.3 Wind along track in the survey area. Arrow vectors show direction and relative speed

CHAPTER 4 RESULTS OF THE ACOUSTIC SURVEY

The distribution area of main groups of pelagic fish in the region, i.e. sardinellas, anchovy, PEL 1 (Clupeids), PEL 2 (mainly carangids) and horse mackerel are depicted in the following figures using the integrator values from the BEI echo-integration system recorded with the ES38B, 38 kHz transducer connected to the EK500. The acoustic densities (in m^2/NM^2) are illustrated by a scale normally used on acoustic surveys with “Dr. Fridtjof Nansen”.

4.1 Guinea Bissau - Liberia


The hydroacoustic survey covered the shelf and slope to 3-500 m bottom depth, with bottom trawls being deployed mainly down to 100 m depth, and continued with one or two pelagic trawls along the transects at night for pelagic species identification. Generally low to medium acoustic densities were found over most of the shelf and only plankton was found in the water column from the shelf break and further offshore. The bottom channel was scrutinized continuously to 500 m bottom depth, but with only few fish targets seen offshore from the shelf break.

Clupeids

Dense concentrations of sardinellas were mainly found on the wide shelf outside of Southeast Guinea Bissau and Guinea, tapering off into Sierra Leone. This coincided well with the upwelling area mentioned above. *Sardinella aurita* dominated in this area (Figure 4.1a).

Further Southeast *Sardinella maderensis* was found, often in the same areas as *S. aurita*. The concentrations were, however, scattered and did not represent significant biomasses. The size of the sardinellas caught was also small. Whether this was due to avoidance of the trawl by the largest fish, or due to a lack of large fish in the area is unclear. However, when large sardinellas are present in a survey area, the “Nansen” trawl normally catches at least some of the large fish, indicating that the wide shelf area mainly contained young sardinella.

Ilisha africana was found in rather loose concentrations in Northwest Guinea Bissau, and on the narrow shelf in Sierra Leone and Liberia, but was absent from the wide shelf in Guinea Bissau, Guinea and Sierra Leone (Figure 4.1b). The total biomass was small.






Figure 4.1. Distribution of sardinellas (a) and P1 (b)(mainly *Ilisha africana*) in the survey area.

Anchovy

Anchovy was found only sparingly in three areas along the coast between 20 and 50 m bottom depth (Figure 4.2). The largest patch was found on the Guinean shelf, with two smaller patches in Sierra Leone and Liberia. The total biomass was small.




Figure 4.2 Distribution of anchovy in the survey area.

*PEL 2 (carangids, scombrids, barracudas and hairtail) and horse mackerel (*Trachurus trecae*)*

The species category PEL 2 consisted of Carangidae, Trichiuridae and Sphyraenidae, and Scombridae. Most pelagic fish were found inshore of 100 m depth. The most abundant PEL 2 species in the trawl catches were *Chloroscombrus chrysurus*, *Trachurus trecae* and *Decapterus punctatus*. Length frequencies of the species can be found in Annex II.

Schools of horse mackerel that could be positively identified were classified as horse mackerel (Figure 4.3b) while suspected but not positively identified horse mackerel schools were included in P2 (Figure 4.3a).

Schools of PEL 2 species, mainly of low density, were found along the whole coastline (Figure 4.3.a) with a high-density patch in northwest Sierra Leonean waters. A substantial part of the biomass was located in the outer part of the wide shelf in the north-western part of the survey area. The total biomass represented the majority of the biomass of pelagic fish in the survey area.

Positively identified horse mackerel (Figure 4.3b) was mainly found in the mid to north-western part of the wide shelf area outside of Guinea and Guinea Bissau, in addition to a small patch in south-eastern Liberian waters. The biomass should be seen in connection with that of PEL 2, and thus represents part of the largest pelagic fish resource in the survey area.




Figure 4.3 Distribution of PEL 2, unidentified carangids, scombrids, barracudas and hairtail (top) and horse mackerel (bottom) in the survey area.

Demersal species

There were consistent acoustic recordings of relatively dense concentrations of demersal fish towards the shelf break. These were mainly *Dentex congensis*, *Dentex angolensis* and *Ariomma bondi*. These sometimes lifted off the shelf, and as a consequence the swept area survey may possibly have underestimated this resource.

4.2 Review of results

No recent previous survey was available for comparisons in the region. Consequently, no comparisons with previous surveys were made. Some general features observed during this survey are commented below.

Pelagic fish were present over large parts of the region. The main densities of pelagic fish were found inshore of 50 m bottom depth, extending inshore into <20 m depth (inshore of the survey area). Most pelagic hauls at night were taken as blind hauls, as relatively few schools were seen on the echo sounder. This was partly due to a dispersed distribution, and partly due to high abundance of plankton that made acoustic detection and separation difficult. Small sardinellas, small carangids and anchovy dominated on the inner shelf, while larger carangids, scombrids and barracudas were more widely distributed over the entire shelf.

The total pelagic fish biomass in the investigated area was small compared to what is normal in upwelling areas like the Canary Current and the Benguela system.

CHAPTER 5 RESULTS FROM THE SWEPT AREA TRAWL SURVEY

The composition of the fish fauna on the continental shelf and slope of the Gulf of Guinea changes with depth (Williams 1968). The catch-distribution analyses were therefore performed for three depth strata on the shelf, 0-50 m (inner shelf) and 51-100 m (outer shelf) and 101-250 m depth (slope). Table 5.1 gives the main species groups with common species in the region. For the different analysis the “other” group includes all species not accounted for in the other groups. Therefore, the content of “other” will change from table to table.

The locations of the trawl stations are shown in Figure 1.1. Records of fishing stations and catches are presented in Annex I and pooled length distributions (weighted by catch) of main species by area are shown in Annex II.

In the swept-area biomass estimates, only the shelf area down to depths of 200 m was included, divided into 0-30 m, 31-50 m 51-100 m and 101-200 m. Mean densities of the main demersal species by depth strata, occurrence and catch distributions are shown in Annex IV.

Table 5.1 Main groups of species included in the analyses of diversity in the Western Gulf of Guinea

Main Groups	Main Families	Typical Species
Demersal	Sciaenidae	<i>Pentheroscion mbizi</i> <i>Pseudotolithus senegalensis</i> <i>Pseudotolithus elongatus</i> <i>Pseudotolithus typus</i>
	Sparidae	<i>Dentex angolensis</i> <i>Dentex congoensis</i> <i>Pagellus bellottii</i> <i>Pagrus caeruleostictus</i> <i>Boops boops</i>
	*Ariidae	<i>Arius latiscutatus</i>
	Serranidae	<i>Serranus accraensis</i> <i>Epinephelus aeneus</i>
	*Lutjanidae	<i>Lutjanus fulgens</i>
	Polynemidae	<i>Galeoides decadactylus</i>
	Haemulidae (=Pomadasyidae)	<i>Brachydeuterus auritus</i> <i>Pomadasys jubelini</i>
	Ophidiidae	<i>Brotula barbata</i>
	*Lethrinidae	<i>Lethrinus atlanticus</i>
	Clupeidae	<i>Sardinella maderensis</i> <i>Sardinella aurita</i> <i>Ilisha africana</i>
Pelagic	Carangidae	<i>Selene dorsalis</i> <i>Chloroscombrus chrysurus</i>

		<i>Decapterus punctatus</i>
		<i>Selar crumenophthalmus</i>
		<i>Caranx hippos</i>
		<i>Caranx cryos</i>
		<i>Alectis alexandrinus</i>
	<i>Scombridae</i>	<i>Scomberomorus tritor</i>
	<i>Trichiuridae</i>	<i>Trichiurus lepturus</i>
	<i>Sphyraenidae</i>	<i>Sphyraena guachancho</i>
Shrimps		<i>Parapenaeus longirostris</i>
		<i>Penaeus notialis</i>
Cephalopods		<i>Sepia officinalis hierredda</i>
		<i>Illex coindetii</i>
		<i>Alloteuthis africana</i>
		<i>Sepiella ornata</i>
		<i>Octopus vulgaris</i>
Sharks and Rays		<i>Raja miraletus</i>
		<i>Squatina oculata</i>
		<i>Mustelus mustelus</i>
Others	<i>Priacanthidae</i>	<i>Priacanthus arenatus</i>
	<i>Citharidae</i>	<i>Citharus linguatula</i>
	<i>Platycephalidae</i>	<i>Grammoplites gruveli</i>
	<i>Synodontidae</i>	<i>Saurida brasiliensis</i>
	<i>Triglidae</i>	<i>Lepidotrigla cadmani</i>
		<i>Lepidotrigla carolae</i>
	<i>Bothidae</i>	<i>Syacium micrurum</i>
	<i>Ariommataidae</i>	<i>Ariomma bondi</i>
	<i>Tetraodontidae</i>	<i>Lagocephalus laevigatus</i>
	<i>Uranoscopidae</i>	<i>Uranoscopus albesca</i>
	<i>Mullidae</i>	<i>Pseudupeneus prayensis</i>
	<i>Fistulariidae</i>	<i>Fistularia petimba</i>
	<i>Cynoglossidae</i>	<i>Cynoglossus canariensis</i>
	<i>Drepanidae</i>	<i>Drepane africana</i>

* Not included in the swept area estimate because of low abundance

5.1 Guinea Bissau

A total of 20 swept-area trawl hauls were made on the Guinea Bissauan shelf. In most hauls the trawl bottom time was around 30 min. This was a prerequisite for the trawl data to be included in the analysis.

Table 5.2 a, b and c shows catch rates by main groups for the inner shelf (0-50 m), mid shelf (51-100 m), outer shelf and slope (101-250 m) shelf, and lower slope respectively. Average catches were around 1053 kg/h on the inner shelf, 181 kg/h on the mid shelf and 1047 kg/h on the outer shelf and slope. The pelagic group contributed 62 % of the total catch and an average catch of 658 kg/h on the inner shelf, while the demersal group accounted for 30 % of

the catch. On the mid shelf, demersal and pelagic species contributed 4 and 60 % or 8 and 108 kg/h respectively while cephalopods accounted for 7 % of the total catch. On the outer shelf and slope the ‘other’ group constituted 63 % of the catch. The demersal group contributed 19%, or 195 kg/h.

Prawns/shrimps were not caught in commercial quantities except for on one station at 229 m.

Sharks and Rays where also present across the shelf in rather low quantities, somewhat more on the mid shelf than shallower and deeper.

The length frequencies of all main species together with the main length – weight parameters are shown in Annex II and III.

Table 5.2 Guinea Bissau. Catch rates (kg/h) by main groups in swept-area bottom-trawl hauls on the a) inner shelf (0-50 m), b) mid shelf (51-100 m) and c) outer shelf and slope (101-250m).

a) Inner shelf, 0-50 m

Sta. no.	Depth	Demersal	Pelagic	Shrimps	Cephalopod	Sharks-rays	Other	Total
987	39	620.0	27.4	2.4	6.2	16.8	25.4	698.1
988	35	429.1	1175.8	0.0	2.2	45.4	45.1	1697.6
992	27	4.9	68.5	0.0	3.0	0.0	27.6	104.0
997	25	1004.0	1371.3	0.0	6.4	18.3	145.3	2545.4
998	36	125.0	194.8	0.0	3.2	0.0	208.8	531.8
1004	28	2.1	896.1	0.0	0.0	0.0	0.0	898.2
1005	43	7.2	870.3	0.0	7.7	0.0	13.2	898.4
Mean	33	313.2	657.7	0.3	4.1	11.5	66.5	1053.4
% catch		29.7	62.4	0.0	0.4	1.1	6.3	100.0

b) Mid shelf, 51-100 m

Sta. no.	Depth	Demersal	Pelagic	Shrimps	Cephalopod	Sharks-rays	Other	Total
989	61	0.0	5.5	0.0	1.3	0.0	12.9	19.8
991	86	31.1	470.0	0.0	56.1	0.0	142.1	699.2
993	51	0.0	6.2	0.0	7.3	0.0	23.8	37.3
994	76	2.4	2.2	0.0	5.6	0.0	54.2	64.4
999	62	11.6	162.8	0.0	6.1	0.0	35.4	215.8
1006	74	0.0	0.0	0.0	0.5	27.2	19.5	47.2
Mean	68	7.5	107.8	0.0	12.8	4.5	48.0	180.6
% catch		4.2	59.7	0.0	7.1	2.5	26.6	100.0

c) Outer shelf and slope, 101-250 m

Sta. no.	Depth	Demersal	Pelagic	Shrimps	Cephalopod	Sharks-rays	Other	Total
990	120	1279.6	0.0	0.0	227.5	0.0	389.1	1896.2
995	165	54.9	29.6	0.0	14.4	12.3	945.6	1056.8

996	246	11.1	0.0	3.4	38.7	0.0	246.5	299.8
1000	142	15.4	164.5	0.0	20.9	0.0	104.3	305.0
1001	229	0.0	301.3	15.2	462.0	4.8	719.5	1502.8
1007	177	0.9	0.0	0.0	1.7	40.2	1903.8	1946.6
1008	220	2.1	5.2	1.1	6.4	16.2	293.9	324.9
Mean	186	194.8	71.5	2.8	110.2	10.5	657.5	1047.4
% catch		18.6	6.8	0.3	10.5	1.0	62.8	100.0

Table 5.3 a, b and c shows the catch rates of the main pelagic families caught in the bottom trawl on the inner, mid and outer shelf/slope respectively. The dominant species on the inner and mid shelf were carangids, represented by *Chloroscombrus chrysurus* and *Chlorophthalmus atlanticus*. Barracudas were sparingly represented in Guinea Bissau, while hairtails (*Trichiurus lepturus*) were only found on 6 stations, with one big catch of 244 kg on the outer shelf dominating the catch.

The clupeoid species that dominates the shallow water pelagic ecosystems in large parts of western Africa were patchily represented on the inner shelf and sparingly represented on the mid shelf. Those that occurred most frequently were *Sardinella aurita* and *Ilisha africana*. In total these species contributed 6.4 % of the total catch on the inner slope.

Table 5.3 Guinea Bissau. Catch rates (kg/h) by main pelagic families in swept-area bottom-trawl hauls on the a) inner shelf (0-50 m), b) mid shelf (51-100 m) and c) outer shelf and slope (101-250).

a) Inner shelf, 0-50 m

Sta. no.	Depth	Clupeoids	Carangids	Scombrids	Hairtails	Barracudas	Other	Total
987	39	1.1	20.8	0.0	0.2	5.4	670.7	698.1
988	35	9.0	1165.4	0.4	0.0	1.0	521.9	1697.6
992	27	0.0	68.5	0.0	0.0	0.0	35.5	104.0
997	25	0.0	1338.3	1.6	0.0	31.5	1174.1	2545.4
998	36	0.1	193.6	1.0	0.0	0.0	337.0	531.8
1004	28	396.0	200.1	300.0	0.0	0.0	2.1	898.2
1005	43	65.0	549.7	255.6	0.0	0.0	28.1	898.4
Mean	33	67.3	505.2	79.8	0.0	5.4	395.6	1053.4
% catch		6.4	48.0	7.6	0.0	0.5	37.6	100.0

b) Mid shelf, 51-100 m

Sta. no.	Depth	Clupeoids	Carangids	Scombrids	Hairtails	Barracudas	Other	Total
989	61	0.0	2.0	0.0	3.5	0.0	14.3	19.8
991	86	0.0	368.0	102.0	0.0	0.0	229.2	699.2
993	51	0.0	6.2	0.0	0.0	0.0	31.1	37.3
994	76	0.0	2.2	0.0	0.0	0.0	62.2	64.4
999	62	24.0	133.0	1.8	3.0	1.0	53.1	215.8

1006	74	0.0	0.0	0.0	0.0	0.0	47.2	47.2
Mean	68	4.0	85.2	17.3	1.1	0.2	72.8	180.6
% catch		2.2	47.2	9.6	0.6	0.1	40.3	100.0

c) Outer shelf and slope, 101-250 m

Sta. no.	Depth	Clupeoids	Carangids	Scombrids	Hairtails	Barracudas	Other	Total
990	120	0.0	0.0	0.0	0.0	0.0	1896.2	1896.2
995	165	0.0	24.0	0.0	5.6	0.0	1027.2	1056.8
996	246	0.0	0.0	0.0	0.0	0.0	299.8	299.8
1000	142	0.4	164.0	0.0	0.0	0.0	140.5	305.0
1001	229	0.0	57.3	0.0	244.0	0.0	1201.5	1502.8
1007	177	0.0	0.0	0.0	0.0	0.0	1946.6	1946.6
1008	220	0.0	0.0	0.0	5.2	0.0	319.7	324.9
Mean	186	0.1	35.1	0.0	36.4	0.0	975.9	1047.4
% catch		0.0	3.3	0.0	3.5	0.0	93.2	100.0

Catch rates of the commercially most important demersal fish groups on the shelf are presented in Table 5.4 a, b and c. The catch rates were in general high on the inner shelf, low on the mid shelf and high again on the outer shelf and slope. Seabreams dominated the inner shelf catches with an average of 154 kg/h, constituting 115% of the total catch in this area. Croakers dominated the outer shelf with 151 kg/h, constituting 14%, one catch in the Northwest part bringing the numbers up.

Snappers, groupers and grunts were only sparingly represented.

Table 5.4 Guinea Bissau. Catch rates (kg/h) of valuable demersal species grouped by families in swept-area bottom-trawl hauls on the a) inner shelf (0-50 m), b) mid shelf (51-100 m) and c) outer shelf and slope (101-250).

a) Inner shelf, 0-50 m

Sta. no.	Depth	Seabreams	Snappers	Groupers	Grunts	Croakers	Other	Total
987	39	6.4	0.0	0.0	0.0	46.1	645.7	698.1
988	35	23.4	0.0	3.6	0.0	0.0	1670.6	1697.6
992	27	2.2	0.0	0.0	2.7	0.0	99.1	104.0
997	25	936.5	0.0	0.0	0.0	0.0	1608.9	2545.4
998	36	110.6	0.0	6.2	8.2	0.0	406.8	531.8
1004	28	0.0	0.0	0.0	0.0	0.0	898.2	898.2
1005	43	0.0	0.0	0.0	0.0	0.0	898.4	898.4
Mean	33	154.1	0.0	1.4	1.6	6.6	889.7	1053.4
% catch		14.6	0.0	0.1	0.1	0.6	84.5	100.0

b) Mid shelf, 51-100 m

Sta. no.	Depth	Seabreams	Snappers	Groupers	Grunts	Croakers	Other	Total
989	61	0	0	0	0	0	19.8	19.8
991	86	31.06	0	0	0	0	668.15	699.2
993	51	0	0	0	0	0	37.28	37.3
994	76	2.4	0	0	0	0	62.01	64.4
999	62	11.62	0	0	0	0	204.22	215.8
1006	74	0	0	0	0	0	47.18	47.2
Mean	68	7.5	0.0	0.0	0.0	0.0	173.1	180.6
% catch		4.2	0.0	0.0	0.0	0.0	95.8	100.0

c) Outer shelf and slope, 101-250 m

Sta. no.	Depth	Seabreams	Snappers	Groupers	Grunts	Croakers	Other	Total
990	120	241.56	0	0	0	1038	616.66	1896.2
995	165	51.22	0	0.28	0	3.38	1001.94	1056.8
996	246	0	0	0	0	1.43	298.35	299.8
1000	142	1.24	0	0	0	14.12	289.62	305.0
1001	229	0	0	0	0	0	1502.76	1502.8
1007	177	0	0	0	0	0	1946.62	1946.6
1008	220	1.8	0	0	0	0	323.06	324.9
Mean	186	42.3	0.0	0.0	0.0	151.0	854.1	1047.4
% catch		4.0	0.0	0.0	0.0	14.4	81.5	100.0

5.2 Guinea

A total of 14 swept-area trawl hauls were made on the shelf off Guinea. The shelf was mainly even, with sandy substrate, and suited for bottom trawling, but with occasional hard spots towards the Southeast.

Table 5.5 a, b and c shows catch rates by main groups for the inner (0-50 m), mid (51-100 m) and outer shelf and slope (101-250 m) respectively. The mean catch rates of pelagic species from 0-50 m depth were 547 kg/h or 61% of the total catch while demersal species contributed 239 kg/h or 27% of the total catch. Prawns/shrimps, cephalopods and sharks and rays contributed only marginally to the total catch in this depth region. The group of other species had a mean catch rate of 100 kg/h or 11% of the total.

The average catch rate at mid shelf was 3711 kg/h. Table 5.5 b shows that pelagic fish was the most abundant group with 92% of the mean total catch. This was mainly due to the contribution from one single catch of pelagic fish. Demersal species had an average catch rate of only 6.8 kg/h.

On the outer shelf and slope cephalopods had the highest catch rates of the main groups with 57.6 kg/h, constituting 8%. “Other” constituted 87%.

Table 5.5 Guinea. Catch rates (kg/h) by main groups in swept-area bottom-trawl hauls on the a) inner shelf (0-50 m), b) mid shelf (51-100 m) and c) outer shelf and slope (101-250).

a) Inner shelf, 0-50 m

Sta. no.	Depth	Demersal	Pelagic	Shrimps	Cephalopod	Sharks-rays	Other	Total
1010	37	13.0	1 062.0	0.0	0.0	0.0	7.5	1 082.5
1011	27	3.4	204.5	0.0	7.2	0.0	97.4	312.4
1020	43	0.7	0.5	0.0	0.9	0.0	23.9	26.1
1021	45	768.3	888.7	0.0	11.1	0.0	33.8	1 701.8
1024	26	392.0	249.5	0.0	5.9	0.0	16.4	663.8
1025	34	136.5	1 421.2	0.0	0.0	0.0	253.8	1 811.5
1026	44	361.0	5.2	0.0	4.3	0.0	270.0	640.5
Mean	37	239.3	547.4	0.0	4.2	0.0	100.4	891.2
% catch		26.8	61.4	0.0	0.5	0.0	11.3	100.0

b) Mid shelf, 51-100 m

Sta. no.	Depth	Demersal	Pelagic	Shrimps	Cephalopod	Sharks-rays	Other	Total
1012	73	0.0	1 091.0	0.0	0.0	0.0	3.0	1 094.0
1018	81	0.0	12 440.6	0.0	1.4	0.0	1 184.6	13 626.5
1019	56	0.3	1.8	0.0	0.8	0.0	21.1	24.0
Mean	70	6.8	4 511.1	0.0	0.7	0.0	402.9	4 914.8
% catch		0.1	91.8	0.0	0.0	0.0	8.2	100.0

c) Outer shelf and slope, 101-300 m

Sta. no.	Depth	Demersal	Pelagic	Shrimps	Cephalopod	Sharks-rays	Other	Total
1013	138	0.0	78.1	0.0	26.0	0.0	318.7	422.9
1014	259	5.9	0.0	2.4	119.0	3.0	137.9	268.2
1016	277	53.9	0.0	2.2	23.1	28.3	500.1	607.6
1017	167	0.0	0.0	0.0	62.4	0.0	1 653.2	1 715.6
Mean	210	15.0	19.5	1.1	57.6	7.8	652.5	753.6
% catch		2.0	2.6	0.2	7.6	1.0	86.6	100.0

The catches of the different pelagic groups in the bottom trawl survey off Guinea is described in Table 5.6. Carangids dominated the pelagic part of the catches on all shelf areas. Catches of carangids comprised 40% on the inner shelf, with an average catch of 358 kg/h. The catches increased to 4500 kg/h on the mid shelf, mainly due to one big catch of *Trachurus trecae*.

Clupeoids had an average catch rate of 179 kg/h on the inner shelf, decreasing to 8 kg/h on the mid shelf. Barracudas were represented on some stations on the inner shelf, but absent on

the mid and outer shelves. Scombrids had catch rates on the mid shelf only of 4 kg/h. The dominating carangids in Guinea were *Trachurus trecae* and *Decapterus punctatus*.

Table 5.6 Guinea. Catch rates (kg/h) by main pelagic families in swept-area bottom-trawl hauls on the a) inner shelf (0-50 m), b) mid shelf (51-100 m), c) outer shelf and slope, 101-250 m.

a) Inner shelf, 0-50 m

Sta. no.	Depth	Clupeoids	Carangids	Scombrids	Hairtails	Barracudas	Other	Total
1010	37	152.9	882.1	0.4	0.0	26.6	20.5	1 082.5
1011	27	8.2	196.3	0.0	0.0	0.0	107.9	312.4
1020	43	0.0	0.5	0.0	0.0	0.0	25.5	26.1
1021	45	85.5	803.2	0.0	0.0	0.0	813.2	1 701.8
1024	26	64.2	173.5	0.0	0.0	11.8	414.3	663.8
1025	34	939.9	448.0	0.0	0.0	33.3	390.3	1 811.5
1026	44	1.3	3.9	0.0	0.0	0.0	635.3	640.5
Mean	37	178.9	358.2	0.1	0.0	10.2	343.9	891.2
% catch		20.1	40.2	0.0	0.0	1.1	38.6	100.0

b) Mid shelf, 51-100 m

Sta. no.	Depth	Clupeoids	Carangids	Scombrids	Hairtails	Barracudas	Other	Total
1012	73	18.8	1 060.6	11.6	0.0	0.0	3.0	1 094.0
1018	81	4.1	12 435.2	1.4	0.0	0.0	1 185.9	13 626.5
1019	56	1.4	0.5	0.0	0.0	0.0	22.2	24.0
Mean	70	8.1	4 498.7	4.3	0.0	0.0	403.7	4 914.8
% catch		0.2	91.5	0.1	0.0	0.0	8.2	100.0

c) Outer shelf and slope, 101-300 m

Sta. no.	Depth	Clupeoids	Carangids	Scombrids	Hairtails	Barracudas	Other	Total
1013	138	0.0	77.0	0.0	1.1	0.0	344.7	422.9
1014	259	0.0	0.0	0.0	0.0	0.0	268.2	268.2
1016	277	0.0	0.0	0.0	0.0	0.0	607.6	607.6
1017	167	0.0	0.0	0.0	0.0	0.0	1 715.6	1 715.6
Mean	210	0.0	19.3	0.0	0.3	0.0	734.0	753.6
% catch		0.0	2.6	0.0	0.0	0.0	97.4	100.0

Catch rates of commercial demersal fish groups in Guinea are presented in Table 5.7 a, b and c. The most dominant group was seabreams with mean catches of 57 kg/h on the inner shelf, and small catches on the mid shelf. The two most important species in this group were *Pagrus caeruleostictus* and *Pagellus bellottii*. Snappers contributed 1% of the total catch on the inner shelf, but were absent further out.

Table 5.7 Guinea. Catch rates (kg/h) of valuable demersal species grouped by families in swept-area bottom-trawl hauls on the a) inner shelf (0-50 m), b) mid shelf (51-100 m) and c) outer shelf and slope (101-250).

a) Inner shelf, 0-50 m

Sta.no.	Depth	Seabreams	Snappers	Groupers	Grunts	Croakers	Other	Total
1010	37	3.1	0.0	0.0	0.0	0.0	1 079.4	1 082.5
1011	27	3.2	0.0	0.0	0.0	0.0	309.2	312.4
1020	43	0.7	0.0	0.0	0.0	0.0	25.3	26.1
1021	45	15.9	0.0	0.0	0.0	0.0	1 685.9	1 701.8
1024	26	79.6	0.0	0.0	0.0	0.0	584.2	663.8
1025	34	108.0	0.0	0.0	0.0	0.0	1 703.5	1 811.5
1026	44	185.9	80.8	13.5	2.5	0.0	357.7	640.5
Mean	37	56.6	11.5	1.9	0.4	0.0	820.8	891.2
% catch		6.4	1.3	0.2	0.0	0.0	92.1	100.0

b) Mid shelf, 51-100 m

Sta.no.	Depth	Seabreams	Snappers	Groupers	Grunts	Croakers	Other	Total
1012	73	0.0	0.0	0.0	0.0	0.0	1 094.0	1 094.0
1018	81	0.0	0.0	0.0	0.0	0.0	13 626.5	13 626.5
1019	56	0.3	0.0	0.0	0.0	0.0	23.7	24.0
Mean	70	0.1	0.0	0.0	0.0	0.0	4 914.7	4 914.8
% catch		0.0	0.0	0.0	0.0	0.0	100.0	100.0

c) Outer shelf and slope, 101-300 m

Sta.no.	Depth	Seabreams	Snappers	Groupers	Grunts	Croakers	Other	Total
1013	138	0.0	0.0	0.0	0.0	0.0	422.9	422.9
1014	259	0.0	0.0	0.0	0.0	0.0	268.2	268.2
1016	277	0.0	0.0	0.0	0.0	0.0	607.6	607.6
1017	167	0.0	0.0	0.0	0.0	0.0	1 715.6	1 715.6
Mean	210	0.0	0.0	0.0	0.0	0.0	753.6	753.6
% catch		0.0	0.0	0.0	0.0	0.0	100.0	100.0

5.3 Sierra Leone

A total of 25 swept-area trawl hauls were made on the shelf off Sierra Leone. The shelf in Sierra Leone is characterised by being wide, with at times hard bottom in the Northwest, narrowing down in the Southeast towards Liberia.

Table 5.8 a, b and c shows catch rates by main groups for the inner (0-50 m), mid (51-100 m) and outer shelf and slope (101-250 m) respectively. The mean catch rates of pelagic species from 0-50 m depth were 471 kg/h or 59% of the total catch while demersal species contributed 203 kg/h or 25% of the total catch. Prawns/shrimps, cephalopods and sharks and

rays contributed somewhat to the total catch with 5.1 kg/h (0.6%), 1.6 kg/h (0.2%) and 15 kg/h (1.9%) respectively on the inner shelf. The group of other species had a mean catch rate of 107 kg/h or 13% of the total on the inner shelf.

The average catch rate on mid shelf was 2014 kg/h. Table 5.5 b shows that pelagic fish was the most abundant group with 25% of the mean total catch. Demersal species had an average catch rate of 135 kg/h, constituting 7%.

On the outer shelf and slope only one haul was carried out due to rough bottom conditions, and the catch at this station was low and mixed.

Table 5.8 Catch rates and related percentages of main groups caught on the inner a) and outer shelf of Sierra Leone (0-50 m), catches in kg/h.

a) Inner shelf, 0-50 m

Sta. no.	Depth	Demersal	Pelagic	Shrimps	Cephalopod	Sharks-rays	Other	Total
1033	39	135.4	16.2	0.0	2.8	0.0	83.3	237.7
1034	30	379.0	154.8	0.0	1.5	0.0	18.6	553.9
1035	26	91.8	30.8	3.8	1.3	0.0	88.8	216.5
1039	32	217.3	653.0	0.0	14.9	0.0	87.5	972.7
1041	32	171.2	1 950.5	0.0	0.0	0.0	151.9	2 273.6
1042	25	786.8	219.2	0.0	0.0	0.0	216.0	1 222.0
1043	26	9.7	91.8	0.0	0.7	0.0	145.7	248.0
1044	26	191.3	2 740.9	0.0	0.0	0.0	115.7	3 047.9
1045	31	12.4	33.4	0.0	3.0	213.2	264.6	526.7
1049	28	192.3	276.8	0.0	0.0	10.5	152.0	631.6
1053	46	386.2	518.0	8.3	0.0	0.0	31.5	944.0
1054	22	192.5	78.6	6.7	0.0	0.6	210.9	489.3
1057	22	90.3	248.3	3.1	0.0	0.0	56.7	398.4
1058	42	234.1	222.5	5.3	1.5	14.5	57.5	535.4
1061	48	59.7	241.1	0.0	0.4	0.0	1.9	303.2
1062	25	104.5	55.0	53.7	0.0	0.0	30.0	243.3
Mean	31	203.4	470.7	5.1	1.6	14.9	107.0	802.8
% catch		25.3	58.6	0.6	0.2	1.9	13.3	100.0

b) Mid shelf, 51-100 m

Sta. no.	Depth	Demersal	Pelagic	Shrimps	Cephalopod	Sharks-rays	Other	Total
1027	70	0.0	28.6	0.0	0.8	0.0	35.1	64.4
1031	78	320.8	451.3	0.0	3.6	6.2	1 901.4	2 683.3
1032	54	0.6	862.8	0.0	0.9	0.0	24.7	889.0
1046	88	42.4	2 214.5	0.0	0.0	0.0	245.3	2 502.3
1051	78	156.8	343.2	0.0	0.0	0.0	79.1	579.1

1052	76	384.9	3.6	0.0	1.3	10.8	20.3	420.7
1059	61	2.7	1.1	0.0	0.0	0.0	5.9	9.8
1060	86	178.3	184.7	0.0	0.0	0.0	8 602.8	8 965.8
Mean	74	135.8	511.2	0.0	0.8	2.1	1 364.3	2 014.3
% catch		6.7	25.4	0.0	0.0	0.1	67.7	100.0

c) Outer shelf and slope, 101-250 m

Sta. no.	Depth	Demersal	Pelagic	Shrimps	Cephalopod	Sharks-rays	Other	Total
1028	108	0.0	3.7	0.0	0.0	0.0	157.4	161.1
Mean	108	0.0	3.7	0.0	0.0	0.0	157.4	161.1
% catch		0.0	2.3	0.0	0.0	0.0	97.7	100.0

The catches of the different pelagic groups in the bottom trawl survey off Sierra Leone is described in Table 5.9. Carangids dominated the pelagic part of the catches on all shelf areas. Catches of carangids comprised 49% on the inner shelf, with an average catch of 386 kg/h. The catches increased to 435 kg/h on the mid shelf. At the one station on the outer shelf only carangids of the main groups were caught, but sparingly (0.2%).

Clupeoids had an average catch rate of 50 kg/h on the inner shelf, increasing to 71 kg/h on the mid shelf. Barracudas were widely and heavily represented on the inner shelf (26 kg/h, 3.3%), but practically absent on the mid and outer shelves. Scombrids had small catch rates on the mid and inner shelf. Hairtails were caught on the inner shelf in the Southeast. The dominating carangids in Sierra Leone were *Chloroscombrus chrysurus*, *Priacanthus arenatus* and *Decapterus punctatus*.

Table 5.9 Sierra Leone. Catch rates (kg/h) by main pelagic families in swept-area bottom-trawl hauls on the a) inner shelf (0-50 m), b) mid shelf (51-100 m), c) outer shelf and slope, 101-250 m.

a) Inner shelf, 0-50 m

Sta. no.	Depth	Clupeoids	Carangids	Scombrids	Hairtails	Barracudas	Other	Total
1033	39	0.0	15.0	0.0	0.0	1.2	221.5	237.7
1034	30	71.6	83.2	0.0	0.0	0.0	399.1	553.9
1035	26	0.8	30.0	0.0	0.0	0.0	185.7	216.5
1039	32	116.0	537.0	0.0	0.0	0.0	319.7	972.7
1041	32	36.0	1 914.5	0.0	0.0	0.0	323.1	2 273.6
1042	25	0.0	219.2	0.0	0.0	0.0	1 002.8	1 222.0
1043	26	0.1	91.7	0.0	0.0	0.0	156.2	248.0
1044	26	0.0	2 739.4	0.0	0.0	1.5	307.0	3 047.9
1045	31	0.0	33.4	0.0	0.0	0.0	493.3	526.7
1049	28	5.2	221.2	3.2	2.6	44.6	354.8	631.6
1053	46	135.4	126.9	6.2	62.0	187.5	426.0	944.0
1054	22	17.2	36.7	3.5	14.4	6.8	410.7	489.3
1057	22	153.9	56.8	0.0	23.6	14.0	150.1	398.4

1058	42	98.1	31.6	1.4	6.4	85.1	312.8	535.4
1061	48	149.2	11.5	0.0	0.0	80.4	62.0	303.2
1062	25	12.5	34.5	0.0	5.9	2.1	188.3	243.3
Mean	31	49.7	386.4	0.9	7.2	26.4	332.1	802.8
% catch		6.2	48.1	0.1	0.9	3.3	41.4	100.0

b) Mid shelf, 51-100 m

Sta. no.	Depth	Clupeoids	Carangids	Scombrids	Hairtails	Barracudas	Other	Total
1027	70	0.5	28.1	0.0	0.0	0.0	35.9	64.4
1031	78	209.2	242.1	0.0	0.0	0.0	2 232.0	2 683.3
1032	54	13.9	848.9	0.0	0.0	0.0	26.3	889.0
1046	88	197.4	1 995.0	22.1	0.0	0.0	287.8	2 502.3
1051	78	0.9	322.9	19.4	0.0	0.0	235.9	579.1
1052	76	0.0	3.6	0.0	0.0	0.0	417.2	420.7
1059	61	0.0	0.1	0.0	0.0	1.0	8.7	9.8
1060	86	147.0	37.7	0.0	0.0	0.0	8 781.1	8 965.8
Mean	74	71.1	434.8	5.2	0.0	0.1	1 503.1	2 014.3
% catch		3.5	21.6	0.3	0.0	0.0	74.6	100.0

c) Outer shelf and slope, 101-250 m

Sta. no.	Depth	Clupeoids	Carangids	Scombrids	Hairtails	Barracudas	Other	Total
1028	108	0.0	3.7	0.0	0.0	0.0	157.4	161.1
Mean	108	0.0	3.7	0.0	0.0	0.0	157.4	161.1
% catch		0.0	2.3	0.0	0.0	0.0	97.7	100.0

Catch rates of commercial demersal fish groups in Sierra Leone are presented in Table 5.10 a, b and c. The most dominant group was seabreams with mean catches of 105 kg/h on the inner shelf, constituting 13 %, and 122 kg/h constituting 6% on the mid shelf. Even in the deep haul at 108 m, 11 kg/h of seabreams were caught. The two most important species in this group were *Pagrus caeruleostictus* and *Pagellus bellottii*, while *Dentex congensis* was caught on the deep station. There was a good catch (22 kg/h) of snappers on one station on the inner shelf. Grunts and croakers were caught in relatively high numbers (18 kg/h, 2.3% and 24 kg/h, 3% respectively) on the inner shelf. Further out they were more or less missing.

Table 5.10 Sierra Leone. Catch rates (kg/h) of valuable demersal species grouped by families in swept-area bottom-trawl hauls on the a) inner shelf (0-50 m), b) mid shelf (51-100 m) and c) outer shelf and slope (101-250).

a) Inner shelf, 0-50 m

Sta. no.	Depth	Seabreams	Snappers	Groupers	Grunts	Croakers	Other	Total
1033	39	135.4	0.0	0.0	0.0	0.0	102.3	237.7
1034	30	129.4	2.4	4.2	241.6	0.0	176.3	553.9
1035	26	90.0	0.0	0.0	0.0	0.0	126.5	216.5

1039	32	170.0	0.0	1.5	0.0	0.0	801.2	972.7
1041	32	163.0	0.0	0.8	0.0	0.0	2 109.7	2 273.6
1042	25	783.2	0.0	0.0	0.0	0.0	438.8	1 222.0
1043	26	9.7	0.0	0.0	0.0	0.0	238.2	248.0
1044	26	176.2	0.0	0.0	12.3	0.0	2 859.4	3 047.9
1045	31	5.8	0.0	0.0	4.8	0.0	516.1	526.7
1049	28	28.3	0.0	0.0	8.2	6.7	588.4	631.6
1053	46	0.0	0.0	1.0	0.0	146.8	796.2	944.0
1054	22	0.0	22.4	0.0	24.9	66.9	375.1	489.3
1057	22	0.0	0.0	0.0	0.4	72.7	325.3	398.4
1058	42	0.0	0.0	0.0	0.0	31.3	504.0	535.4
1061	48	3.8	0.0	8.9	0.0	1.7	288.7	303.2
1062	25	0.0	0.0	0.0	0.0	60.2	183.1	243.3
Mean	31	105.9	1.6	1.0	18.3	24.2	651.8	802.8
% catch		13.2	0.2	0.1	2.3	3.0	81.2	100.0

b) Mid shelf, 51-100 m

Sta. no.	Depth	Seabreams	Snappers	Groupers	Grunts	Croakers	Other	Total
1027	70	0.0	0.0	0.0	0.0	0.0	64.4	64.4
1031	78	280.0	0.0	13.8	0.0	0.0	2 389.5	2 683.3
1032	54	0.6	0.0	0.0	0.0	0.0	888.4	889.0
1046	88	39.7	0.0	0.0	0.0	0.0	2 462.6	2 502.3
1051	78	128.3	0.0	21.6	0.0	0.0	429.1	579.1
1052	76	378.4	1.0	0.0	0.0	5.5	35.9	420.7
1059	61	1.8	0.0	0.0	0.0	0.0	8.0	9.8
1060	86	150.8	0.0	0.0	0.0	0.0	8 815.0	8 965.8
Mean	74	122.4	0.1	4.4	0.0	0.7	1 886.6	2 014.3
% catch		6.1	0.0	0.2	0.0	0.0	93.7	100.0

c) Outer shelf and slope, 101-250 m

Sta. no.	Depth	Seabreams	Snappers	Groupers	Grunts	Croakers	Other	Total
1028	108	10.8	0.0	0.0	0.0	0.0	150.3	161.1
Mean	108	10.8	0.0	0.0	0.0	0.0	150.3	161.1
% catch		0.5	0.0	0.0	0.0	0.0	6.7	100.0

5.4 Liberia

The cost of Liberia is generally characterised by a narrow shelf that breaks at around 100 m depth. A total of 20 swept-area trawl hauls were carried out. Trawling was made difficult towards the shelf break and shore due to rough bottom conditions, particularly in the Southeast.

Table 5.11 a, b, c and d shows catch rates by main groups for the inner (0-50 m) and mid (51-100 m) shelf respectively, as no trawling was possible in the outer shelf region. Pelagic species dominated in the inshore region with 162 kg/h or 52% of the catches. The second most important group was demersal species which contributed 95 kg/h and 30% of the catches. Prawns/shrimps contributed 19 kg/h (6.2%) while cephalopods and sharks only contributed smaller amounts. The group of other species had a mean catch rate of 36 kg/h or 11% of the total. Overall catches were about the same on the inner and mid shelf with 314 and 320 kg/h respectively. On the mid shelf demersal species dominated with 144 kg/h and 45%, followed by pelagics with 91 kg/h and 28%. The group of other species gave 71 kg/h and 22% of the catches. Prawns and shrimps, cephalopods and sharks and rays only contributed small amounts to the total catches.

Table 5.11 Liberia. Catch rates (kg/h) by main groups in swept-area bottom-trawl hauls on the a) inner shelf (0-50 m), b) mid shelf (51-100 m) and c).

a) Inner shelf, 0-50 m

Sta. no.	Depth	Demersal	Pelagic	Shrimps	Cephalopod	Sharks-rays	Other	Total
1065	31	194.8	327.9	46.3	0.0	0.0	37.3	606.3
1069	47	74.5	111.9	5.7	2.4	0.0	8.1	202.6
1070	25	117.0	96.8	32.6	0.0	0.0	18.7	265.2
1073	25	4.4	15.8	1.0	0.0	0.0	29.5	50.7
1078	44	111.3	9.9	29.8	1.8	0.0	48.7	201.5
1087	39	69.5	407.1	0.3	0.0	2.3	76.2	555.4
1088	30	1 013.4	33.1	0.6	0.0	0.0	34.7	1 081.8
Mean	34	226.4	143.2	16.6	0.6	0.3	36.2	423.4
% catch		53.5	33.8	3.9	0.1	0.1	8.5	100.0

b) Mid shelf, 51-100 m

Sta. no.	Depth	Demersal	Pelagic	Shrimps	Cephalopod	Sharks-rays	Other	Total
1066	56	77.8	5.7	0.5	0.1	0.0	0.9	85.1
1067	80	84.9	3.4	0.4	7.2	12.0	4.4	112.2
1068	94	57.0	0.2	0.0	6.7	2.8	9.7	76.3
1074	61	21.8	44.8	0.0	9.7	0.0	4.1	80.4
1075	94	246.1	105.0	0.0	0.0	0.0	75.5	426.6
1076	90	230.6	0.0	0.0	0.6	0.0	19.4	250.6
1077	64	156.3	8.2	2.5	2.2	0.0	46.7	215.9
1080	59	49.0	27.9	0.1	4.8	0.0	6.4	88.1
1081	72	75.2	77.3	0.0	3.9	0.0	13.8	170.3
1082	89	618.1	537.8	0.0	2.7	31.7	91.8	1 282.1
1083	92	117.8	66.5	0.0	1.2	90.0	15.1	290.6
1084	71	106.8	3.4	0.0	6.1	1.4	42.6	160.2
1085	80	160.4	276.4	0.0	1.8	6.3	647.2	1 092.1
1086	79	14.9	111.5	0.1	6.2	0.0	13.3	146.0
Mean	77	144.0	90.6	0.3	3.8	10.3	70.8	319.7

% catch	45.0	28.3	0.1	1.2	3.2	22.1	100.0
---------	------	------	-----	-----	-----	------	-------

Pelagic species were fairly abundant on the Liberian shelf. Clupeoids dominated the inner shelf (48 kg/h, 15%), while Carangids dominated the mid shelf (65 kg/h, 20%). The dominant clupeids were *Sardinella maderensis* and *Ilisha africana*. The carangids were dominated by *Selene dorsalis* and *Decapterus punctatus*.

Table 5.12 Liberia. Catch rates (kg/h) by main pelagic families in swept-area bottom-trawl hauls on the a) inner shelf (0-50 m), b) mid shelf (51-100 m).

a) Inner shelf, 0-50 m

Sta. no.	Depth	Clupeoids	Carangids	Scombrids	Hairtails	Barracudas	Other	Total
1065	31	198.6	25.2	0.0	42.4	61.7	278.4	606.3
1069	47	65.9	12.4	0.0	8.8	24.8	90.7	202.6
1070	25	18.5	48.0	0.0	20.9	9.5	168.4	265.2
1073	25	0.2	0.1	1.2	0.0	14.3	34.9	50.7
1078	44	2.3	1.4	0.0	1.5	4.7	191.7	201.5
1087	39	4.7	59.4	0.6	0.7	341.8	148.2	555.4
1088	30	0.0	33.1	0.0	0.0	0.0	1048.7	1081.8
Mean	34	41.5	25.7	0.3	10.6	65.3	280.1	423.4
% catch		9.8	6.1	0.1	2.5	15.4	66.2	100.0

b) Mid shelf, 51-100 m

Sta. no.	Depth	Clupeoids	Carangids	Scombrids	Hairtails	Barracudas	Other	Total
1066	56	0.6	1.3	0.0	0.8	2.9	79.3	85.1
1067	80	0.0	0.5	0.0	2.9	0.0	108.8	112.2
1068	94	0.0	0.0	0.0	0.2	0.0	76.1	76.3
1074	61	0.2	43.0	0.0	0.0	1.7	35.5	80.4
1075	94	0.0	105.0	0.0	0.0	0.0	321.6	426.6
1076	90	0.0	0.0	0.0	0.0	0.0	250.6	250.6
1077	64	0.3	1.0	0.0	0.0	6.9	207.7	215.9
1080	59	0.6	26.1	0.0	0.0	1.2	60.3	88.1
1081	72	22.0	55.3	0.0	0.0	0.0	93.0	170.3
1082	89	178.2	352.4	7.2	0.0	0.0	744.3	1 282.1
1083	92	0.0	63.0	3.5	0.0	0.0	224.1	290.6
1084	71	0.1	2.5	0.0	0.0	0.9	156.8	160.2
1085	80	2.2	236.3	6.1	0.0	31.8	815.7	1 092.1
1086	79	2.7	17.3	0.0	3.5	88.0	34.5	146.0
Mean	77	14.8	64.5	1.2	0.5	9.5	229.2	319.7
% catch		4.6	20.2	0.4	0.2	3.0	71.7	100.0

On the inner shelf no seabreams were caught. Grunts were caught on most stations, while the dominating group was croakers, with 67 kg/h and 21% of the mean catches. “Others” constituted 238 kg/h and 76% of the mean catches. Sciaenids were prominent on the innermost station on the last transect, as it was trawled just outside of an estuary.

On the mid shelf the seabreams dominated, with a mean catch rate of 100 kg/h and 31% of the mean total catch. The main species were *Dentex congoensis* and *D. angolensis*. Croakers were also caught here, but in much smaller quantities. “Others” constituted 212 kg/h and 66%.

Table 5.13 Liberia. Catch rates (kg/h) of valuable demersal species grouped by families in swept-area bottom-trawl hauls on the a) inner shelf (0-50 m), b) mid shelf (51-100 m) and c) outer shelf and slope (101-250).

a) Inner shelf, 0-50 m

Sta.no.	Depth	Seabreams	Snappers	Groupers	Grunts	Croakers	Other	Total
1065	31	0.0	0.0	0.0	0.3	182.1	423.9	606.3
1069	47	0.0	0.0	0.0	0.0	15.3	187.4	202.6
1070	25	0.0	0.0	0.0	2.5	97.4	165.3	265.2
1073	25	0.0	0.0	0.0	0.5	3.9	46.3	50.7
1078	44	0.0	0.0	0.0	47.5	51.3	102.7	201.5
1087	39	0.0	0.0	0.0	1.0	51.9	502.4	555.4
1088	30	0.0	46.1	0.0	1.4	917.9	116.4	1 081.8
Mean	34	0.0	6.6	0.0	7.6	188.6	220.6	423.4
% catch		0.0	1.6	0.0	1.8	44.5	52.1	100.0

b) Mid shelf, 51-100 m

Sta.no.	Depth	Seabreams	Snappers	Groupers	Grunts	Croakers	Other	Total
1066	56	1.2	0.0	17.6	0.0	0.7	65.6	85.1
1067	80	72.6	0.0	12.1	0.0	0.0	27.5	112.2
1068	94	51.6	0.0	0.2	0.0	5.1	19.3	76.3
1074	61	16.5	0.0	2.3	0.6	0.0	61.0	80.4
1075	94	189.9	0.0	0.0	0.0	0.0	236.7	426.6
1076	90	229.6	0.0	0.0	0.0	0.0	20.9	250.6
1077	64	97.0	0.0	0.0	0.0	2.4	116.6	215.9
1080	59	33.4	0.0	13.9	0.0	0.8	40.0	88.1
1081	72	75.2	0.0	0.0	0.0	0.0	95.0	170.3
1082	89	392.2	0.0	0.0	0.0	23.4	866.5	1 282.1
1083	92	92.5	0.0	0.0	0.0	1.2	196.9	290.6
1084	71	77.1	0.0	15.8	0.0	2.2	65.2	160.2
1085	80	66.0	0.0	0.0	0.0	3.0	1 023.1	1 092.1
1086	79	3.2	0.0	0.0	0.0	11.7	131.1	146.0
Mean	77	99.8	0.0	4.4	0.0	3.6	211.8	319.7
% catch		31.2	0.0	1.4	0.0	1.1	66.2	100.0

5.5 Swept area estimates

Swept area estimates based on the demersal trawl data are presented in Table 5.14, and in further detail in Appendix VIII.

Table 5.14 Total swept area estimates in tonnes 2006

Country	Seabreams	Grunts ¹	Croakers	Groupers	Snappers	<i>B. auritus</i>	Sharks	Rays	Cephalop.	Total
Guinea Bissau	15907	142	11736	128	0	11622	1809	1865	3837	47047
Guinea	7873	41	0	287	1804	19823	189	358	1932	32306
Sierra Leone	21362	2131	2897	440	191	5940	2144	298	260	35663
Liberia	13031	369	9535	440	297	1628	1056	306	407	27068
Total	58173	2683	24168	1295	2292	39013	5198	2826	6435	142084

¹ Grunts excluding *Brachydeuterus auritus*.

The total swept area biomass for the shelf of the four countries in question was measured to be around 140,000 tonnes. A number of uncertainties should be considered when applying this and other abundance estimates in this report, particularly regarding trawl catchability, patchiness, interpolation and area calculations. Also, the relatively low number of trawl hauls per area unit, relative to the degree of variability in fish concentrations should be remembered. So, this being said, the numbers can be used as a guideline to magnitudes regarding the resource situation in the area.

Guinea Bissau had the highest abundance, constituted by good availability of seabreams, croakers and cephalopods, pluss the ever-present *Brachydeuterus auritus*. Also, the size structure in the populations seemed to be good, indicating healthy fish stocks in terms of exploitation. The seasonal upwelling situation probably contributes towards this situation, but looking at the stock structure, there seems to be an indication of a balanced fishing effort, at least on the southern shelf that was the object of investigation during this survey.

Guinea also had a fairly large demersal fish resource constituted by seabreams and quite a few snappers in addition to large amounts of *Brachydeuterus*, but given the wide shelf and the influence of the upwelling, more fish could have been expected. Also, the length structure in the populations was not as good as in Guinea Bissau, indicating high fishing pressure. This was emphasised by the observation of large international trawlers fishing particularly in the shelf break area. Indications of good recruitment was, however, present, as small commercial fish were abundantly present.

In Sierra Leone the situation improved again. The total abundance measured was comparable to Guinea, but the fish were larger, and consequently the fishing pressure seemed to be lower than on the Guinean shelf. Seabreams constituted the most frequent catch of valuable commercial species.

Liberia had the lowest measured abundance, but not substantially lower than Sierra Leone and Guinea. Also here, seabreams were the dominating group, followed by croakers in terms of valuable commercial species. It should, however, be noted that one big catch of large croakers close to a probable spawning habitat did raise these numbers somewhat. The exploitation situation in Liberia seemed good, with accumulated stock type length distributions being the norm.

5.6 Overall review of results

Guinea Bissau

The survey off Guinea Bissau was characterised by fairly good catches of both demersal and pelagic fish. This apparent comparative richness probably originates from the seasonal upwelling, creating the nutrient basis for a relatively high primary production for this latitude. The demersal fish generally had a quite large mean length for the species in question, indicating that the fishing pressure is moderate.

Guinea

Guinea has a wide shelf that is also exposed to the seasonal upwelling. A high fish production for the latitude should therefore be expected. The preliminary results of the survey may indicate that the fishing pressure is quite high, as the mean lengths of both the demersal and pelagic fishes are quite low for the species in question. There was, however, no clear indication of a recruitment failure, as juvenile fish were abundant.

Sierra Leone

Sierra Leone gave good catches both of pelagic and particularly of large demersal fish. This indicates that the fishing pressure here has been lower than in Guinea, as the upwelling effect should be less here than further to the Northwest. While the present situation is good, it may soon change, as a fleet of IUU fishing vessels were operating in the region.

Liberia

Liberia is characterised by an oceanic environment with higher sea surface temperatures than Guinea Bissau, Guinea and Northwestern Sierra Leone, at least during the upwelling in the rainy season. The demersal species dominated, but also pelagic species like sardinella were found. There seemed to have been a low fishing pressure particularly in the Southeastern part, as large fish were abundant close to shore. Liberia has tropical shelf waters, and consequently can expect low production. There are, however, still potentially good conditions for artisanal fishing and sport fishing, as much of the shelf area is rocky, and thus is protected from illegal trawl fishing.

Annex I Records of fishing stations

PROJECT STATION: 987						BDEPTH:	120	120	Validity code:
						Towing dir:	83°	Wire out:	360 m Speed: 30 kn*10
DATE:29/ 4/06	GEAR TYPE: BT No:19	POSITION:Lat	N	1149	Long	W	1709		
start	stop	duration							
TIME :21:04:06	21:28:09	24	(min)	Purpose code: 3					
LOG :6509.18	6510.51	1.32		Area code : 1					
FDEPTH:	39	38		GearCond.code:					
BDEPTH:	39	38		Validity code:					
Towing dir: 170° Wire out: 150 m Speed: 30 kn*10									
Sorted: 119 Kg	Total catch:	279.23	CATCH/HOUR:	698.08					
SPECIES	CATCH/HOUR	% OF TOT. C	SAMP						
	weight	numbers							
Brachydeuterus auritus	456.85	5460	65.44	4277	Pentheroscion mbizi	1015.20	10672	53.54	4290
Arius parkii	110.68	348	15.85	4278	Spicara alta	297.60	1740	15.69	4293
Pseudotolithus senegalensis	31.00	30	4.44	4273	Todaropsis eblanae	222.00	3664	11.71	4294
Chloroscombrus chrysurus	20.05	270	2.87	4275	Dentex congogensis	116.40	842	6.14	4291
Galeoides decadactylus	18.13	35	2.60	4274	Dentex angolensis	79.80	528	4.21	4288
Rhizoprionodon acutus	16.75	3	2.40		Scorpaena stephanica	60.00	180	3.16	
Pteroscion peli	12.10	138	1.73	4276	Pagrus africanus	31.80	72	1.68	4287
Pagellus bellottii	6.35	28	0.91		Umbrina canariensis	22.80	72	1.20	
Sepiella ornata	6.15	55	0.88		Dentex macrophthalmus	13.56	60	0.72	4292
Sphyraena guachancho	5.40	8	0.77		Zeus faber	11.00	20	0.58	4289
Syacium micrurum	3.85	250	0.55		Illlex coindetii	5.52	36	0.29	
Pseudotolithus typus	3.00	5	0.43		Priacanthus arenatus	5.40	24	0.28	
Penaeus notialis	2.38	125	0.34		Aulopus cadenati	3.48	36	0.18	
Chilomycterus spinosus mauret.	2.15	8	0.31		Zenopsis conchifer	3.00	2	0.16	
Sardinella maderensis	0.98	8	0.14		Chaetodon hoefleri	2.88	24	0.15	
Pseudupeneus prayensis	0.70	8	0.10		Antigonion capros	2.76	48	0.15	
Selene dorsalis	0.70	8	0.10		Ariommha bondi	1.56	36	0.08	
Eucinostomus melanopterus	0.43	8	0.06		Sphoeroides pachgaster	1.46	2	0.08	
OPICHTHIDAE	0.15	15	0.02		Total	1896.22		100.00	
Trichiurus lepturus	0.15	15	0.02						
Ilisha africana	0.15	8	0.02						
Total	698.10	99.98							
PROJECT STATION: 988									
DATE:30/ 4/06	GEAR TYPE: BT No:19	POSITION:Lat	N	1115	Long	W	1703		
start	stop	duration							
TIME :07:02:40	07:32:43	30	(min)	Purpose code: 3					
LOG :6585.73	6587.18	1.44		Area code : 1					
FDEPTH:	32	38		GearCond.code:					
BDEPTH:	32	38		Validity code:					
Towing dir: 270° Wire out: 150 m Speed: 30 kn*10									
Sorted: 160 Kg	Total catch:	848.82	CATCH/HOUR:	1697.64					
SPECIES	CATCH/HOUR	% OF TOT. C	SAMP						
	weight	numbers							
Trachurus trecae	671.20	10662	39.54	4285	SPECIES	CATCH/HOUR	% OF TOT. C	SAMP	
Brachydeuterus auritus	381.20	10334	22.45	4287		weight	numbers		
Decapterus rhonchus	376.20	1652	22.16	4283	Trachurus trecae	241.33	3257	34.52	4296
Chloroscombrus chrysurus	95.00	988	5.60	4284	Decapterus rhonchus	126.67	273	18.12	4295
Rhizoprionodon acutus	45.40	34	2.67		Scomber japonicus	102.00	2073	14.59	4299
Pseudupeneus prayensis	29.88	250	1.76	4280	Ariommha bondi	66.67	977	9.54	4297
Pagellus bellottii	23.38	176	1.38	4279	Todaropsis eblanae	47.33	1063	6.77	
Decapterus punctatus	23.00	850	1.35	4286	Scorpaena stephanica	36.00	120	5.15	
Arius parkii	14.50	38	0.85		Dentex congogensis	30.13	460	4.31	4298
Sardinella aurita	9.00	288	0.53	4281	Raja miraletus	11.87	20	1.70	
Arius latiscutatus	6.40	2	0.38		Antigonion capros	9.67	153	1.38	
Fistularia tabacaria	5.60	6	0.33		Illex coindetii	8.27	140	1.18	
Lagocephalus laevigatus	5.12	12	0.30		Sphoeroides pachgaster	7.33	20	1.05	
Epinephelus aeneus	3.62	12	0.21	4282	Zeus faber	5.67	13	0.81	
Syacium micrurum	3.26	12	0.19		Chaetodon hoefleri	2.33	13	0.33	
Alloteuthis africana	1.12	238	0.07		Dactylopterus volitans	1.67	7	0.24	
Sepiella ornata	1.12	26	0.07		Dentex angolensis	0.93	7	0.13	
Sphyraena guachancho	1.00	2	0.06		Spicara alta	0.67	7	0.10	
Sphoeroides marmoratus	0.76	12	0.04		Sepiella ornata	0.47	7	0.07	
Bothus podas africanus	0.50	12	0.03		Lepidotrigla carolae	0.13	7	0.02	
Scomber japonicus	0.38	26	0.02		Aulopus cadenati	0.07	7	0.01	
Total	1697.64	99.99			Total	699.21		100.02	
PROJECT STATION: 989									
DATE:30/ 4/06	GEAR TYPE: BT No:19	POSITION:Lat	N	1117	Long	W	1711		
start	stop	duration							
TIME :10:11:46	10:35:20	24	(min)	Purpose code: 3					
LOG :6601.29	6602.50	1.21		Area code : 1					
FDEPTH:	61	60		GearCond.code:					
BDEPTH:	61	60		Validity code:					
Towing dir: 140° Wire out: 200 m Speed: 30 kn*10									
Sorted: 8 Kg	Total catch:	7.90	CATCH/HOUR:	19.75					
SPECIES	CATCH/HOUR	% OF TOT. C	SAMP						
	weight	numbers							
Fistularia petimba	5.93	263	30.03		SPECIES	CATCH/HOUR	% OF TOT. C	SAMP	
Trichiurus lepturus	3.50	10	17.72			weight	numbers		
Fistularia petimba	3.13	8	15.85		Decapterus rhonchus	64.00	210	61.54	4301
Caranx crysos	2.03	3	10.28		Balistes capriscus	17.00	18	16.35	4302
Liocarcarinus corrugatus	1.68	385	8.51		Pomadasys rogeri	2.72	2	2.62	
Lagocephalus laevigatus	1.00	3	5.06		Sepia officinalis hierredda	2.64	10	2.54	
Alloteuthis africana	0.73	263	3.70		Rhinobatos rhinobatos	2.34	2	2.25	
Sepiella ornata	0.50	5	2.53		Pagellus bellottii	2.16	12	2.08	4300
Chelidonichthys gabonensis	0.33	3	1.67		Chloroscombrus chrysurus	1.96	14	1.88	
Dicologlossa cuneata	0.28	3	1.42		Echeneis naucrates	1.86	4	1.79	
Priacanthus arenatus	0.23	3	1.16		Trachurus trecae	1.68	10	1.62	
Saurida brasiliensis	0.18	15	0.91		Lagocephalus laevigatus	1.56	2	1.50	
Sepia officinalis hierredda	0.10	3	0.51		Diomedea sp.	1.22	2	1.17	
Microchirius boscanion	0.05	5	0.25		Rachycentron canadum	1.04	2	1.00	
Gammoplites gruvelli	0.05	3	0.25		Chilomycterus spinosus mauret.	0.90	2	0.87	
Lepidotrigla carolae	0.05	3	0.25		Selene dorsalis	0.70	6	0.67	
Syacium micrurum	0.03	3	0.15		Chaetodon hoefleri	0.48	4	0.46	
Total	19.80	100.25			Uranoscopus polli	0.42	2	0.40	
PROJECT STATION: 990									
DATE:30/ 4/06	GEAR TYPE: BT No:19	POSITION:Lat	N	1115	Long	W	1719		
start	stop	duration							
TIME :12:46:45	13:16:45	30	(min)	Purpose code: 3					
LOG :6620.90	6622.40	1.50		Area code : 1					
FDEPTH:	120	120		GearCond.code:					
BDEPTH:	50	52		Validity code:					
Towing dir: 140° Wire out: 200 m Speed: 30 kn*10									
Sorted: 19 Kg	Total catch:	19.26	CATCH/HOUR:	37.28					
SPECIES	CATCH/HOUR	% OF TOT. C	SAMP						
	weight	numbers							
Penheroscion mbizi	1015.20	10672	53.54						
Spicara alta	297.60	1740	15.69						
Todaropsis eblanae	222.00	3664	11.71						
Dentex congogensis	116.40	842	6.14						
Dentex angolensis	79.80	528	4.21						
Scorpaena stephanica	60.00	180	3.16						
Pagrus africanus	31.80	72	1.68						
Umbrina canariensis	22.80	72	1.20						
Dentex macrophthalmus	13.56	60	0.72						
Zeus faber	11.00	20	0.58						
Illex coindetii	5.52	36	0.29						
Priacanthus arenatus	5.40	24	0.28						
Aulopus cadenati	3.48	36	0.18						
Zenopsis conchifer	3.00	2	0.16						
Chaetodon hoefleri	2.88	24	0.15						
Antigonion capros	2.76	48	0.15						
Ariommha bondi	1.56	36	0.08						
Sphoeroides pachgaster	1.46	2	0.08						
Total	1896.22								
PROJECT STATION: 991									
DATE:30/ 4/06	GEAR TYPE: BT No:19	POSITION:Lat	N	1115	Long	W	1717		
start	stop	duration							
TIME :15:51:10	16:09:24	18	(min)	Purpose code: 3					
LOG :6641.39	6642.32	0.92		Area code : 1					
FDEPTH:	86	85		GearCond.code:					
BDEPTH:	86	85		Validity code:					
Towing dir: 90° Wire out: 236 m Speed: 30 kn*10									
Sorted: 105 Kg	Total catch:	209.76	CATCH/HOUR:	699.20					
SPECIES	CATCH/HOUR	% OF TOT. C	SAMP						
	weight	numbers							
Trachurus trecae	241.33	3257	34.52						
Decapterus rhonchus	126.67	273	18.12						
Scomber japonicus	102.00	2073	14.59						
Ariommha bondi	66.67	977	9.54						
Todaropsis eblanae	47.33	1063	6.77						
Scorpaena stephanica	36.00	120	5.15						
Dentex congogensis	30.13	460	4.31						
Raja miraletus	11.87	20	1.70						
Antigonion capros	9.67	153	1.38						
Illex coindetii	8.27	140	1.18						
Sphoeroides pachgaster	7.33	20	1.05						
Zeus faber	5.67	13	0.81						
Chaetodon hoefleri	2.33	13	0.33						
Dactylopterus volitans	1.67	7	0.24						
Dentex angolensis	0.93	7	0.13						
Spicara alta	0.67	7	0.10						
Sepiella ornata	0.47	7	0.07						
Lepidotrigla carolae	0.13	7	0.02						
Aulopus cadenati	0.07	7	0.01						
Total	699.21								
PROJECT STATION: 992									
DATE: 1/ 5/06	GEAR TYPE: BT No:19	POSITION:Lat	N	1104	Long	W	1653		
start	stop								

	weight	numbers		
Fistularia petimba	13.55	58	36.35	
Sepia officinalis hierredda	7.28	35	19.53	4303
Lagocephalus laevigatus	5.23	10	14.03	
Caranx cryos	2.71	4	7.27	
Fistularia tabacaria	2.61	2	7.00	
Decapterus rhonchus	1.94	8	5.20	
Chaetodon hoefleri	0.87	6	2.33	
Decapterus punctatus	0.79	8	2.12	
Trachurus trecae	0.77	4	2.07	
Chelidonichthys gabonensis	0.74	6	1.98	
Sphoeroides marmoratus	0.33	4	0.89	
Scorpaena scrofa	0.21	2	0.56	
Syacium micrurum	0.15	2	0.40	
Bothus podas africanus	0.10	2	0.27	
Total	37.28	100.00		

Lepidotrigla carolae	0.66	31	0.22
Echelus myrus	0.58	2	0.19
Synchiropus phaeton	0.39	21	0.13
Peristedion cataphractum	0.35	17	0.12
Coelorinchus coelorrhincus	0.31	4	0.10
Grammoplites griseus	0.31	4	0.10
Chascanopsetta lugubris	0.21	4	0.07
Cynoponticus ferox	0.14	4	0.05
Zenion longipinnis	0.04	17	0.01
Total	299.78	100.02	

PROJECT STATION: 994
DATE: 1/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 1059
start stop duration Long W 1704
TIME :11:46:02 12:16:39 31 (min) Purpose code: 3
LOG :6749.90 6751.34 1.42 Area code : 1
FDEPTH: 76 76 GearCond.code:
BDEPTH: 76 76 Validity code:
Towing dir: 135° Wire out: 240 m Speed: 30 kn*10
Sorted: 33 Kg Total catch: 33.27 CATCH/HOUR: 64.39

PROJECT STATION: 997
DATE: 2/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 1044
start stop duration Long W 1625
TIME :09:33:35 10:08:56 35 (min) Purpose code: 3
LOG :6896.65 6898.39 1.72 Area code : 1
FDEPTH: 23 26 GearCond.code:
BDEPTH: 23 26 Validity code:
Towing dir: 245° Wire out: 150 m Speed: 30 kn*10

Sorted: 193 Kg Total catch: 1484.83 CATCH/HOUR: 2545.42
SPECIES CATCH/HOUR % OF TOT. C SAMP
weight numbers
Fistularia petimba 16.65 33 25.86
Pseudupeneus prayensis 10.41 72 16.17 4305
Scorpaena stephanica 6.70 21 10.41
Octopus vulgaris 4.26 2 6.62
Raja miraletus 4.24 10 6.58
Dactylopterus volitans 3.35 8 5.20
Sphoeroides pacchaster 2.77 4 4.30
Chaetodon hoefleri 2.75 19 4.27
Pagrus africanus 2.09 4 3.25 4304
Uranoscopus albusca 1.84 2 2.86
Chelidonichthys gabonensis 1.72 14 2.67
Priacanthus arenatus 1.43 17 2.22
Sepia officinalis hierredda 1.28 10 1.99
Echelus myrus 1.26 2 1.96
Caranx cryos 0.79 2 1.23
Torpedo torpedo 0.79 4 1.23
Decapterus rhonchus 0.74 2 1.15
Decapterus punctatus 0.64 6 0.99
Pagellus bellottii 0.31 14 0.48
Fistularia petimba 0.29 31 0.45
Alloteuthis africana 0.10 23 0.16
Total 64.41 100.05

PROJECT STATION: 994
DATE: 1/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 1059
start stop duration Long W 1704
TIME :11:46:02 12:16:39 31 (min) Purpose code: 3
LOG :6749.90 6751.34 1.42 Area code : 1
FDEPTH: 76 76 GearCond.code:
BDEPTH: 76 76 Validity code:
Towing dir: 135° Wire out: 240 m Speed: 30 kn*10
Sorted: 33 Kg Total catch: 1484.83 CATCH/HOUR: 2545.42
SPECIES CATCH/HOUR % OF TOT. C SAMP
weight numbers
Chloroscombrus chrysurus 1039.89 18329 40.85 4317
Pagellus bellottii 772.97 4666 30.37 4315
Decapterus rhonchus 253.03 2745 9.94 4319
Pagrus caeruleostictus 163.54 602 6.42 4318
Pseudupeneus prayensis 101.06 864 3.97 4316
Brachydeuterus auritus 67.53 722 2.65 4320
Albula vulpes 35.64 46 1.40
Caranx cryos 28.70 93 1.13
Sphyraena afra 27.60 2 1.08
Rhizoprionodon acutus 18.34 5 0.72
Selene dorsalis 16.66 170 0.65
Sphyraena guachancho 3.86 15 0.15
Nicholsina usta 3.86 15 0.15
Sepia officinalis hierredda 3.70 15 0.15
Ephippion guttifer 2.74 3 0.11
Octopus vulgaris 2.74 3 0.11
Panulirus regius 1.97 2 0.08
Scomberomorus tritor 1.59 2 0.06
Total 2545.42 99.99

PROJECT STATION: 995
DATE: 1/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 1054
start stop duration Long W 1712
TIME :14:10:31 14:40:58 30 (min) Purpose code: 3
LOG :6763.82 6765.35 1.54 Area code : 1
FDEPTH: 178 152 GearCond.code:
BDEPTH: 178 152 Validity code:
Towing dir: 62° Wire out: 470 m Speed: 30 kn*10
Sorted: 141 Kg Total catch: 528.47 CATCH/HOUR: 1056.94

PROJECT STATION: 998
DATE: 2/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 1042
start stop duration Long W 1629
TIME :11:00:30 11:30:38 30 (min) Purpose code: 3
LOG :6901.60 6903.17 1.56 Area code : 1
FDEPTH: 35 37 GearCond.code:
BDEPTH: 35 37 Validity code:
Towing dir: 245° Wire out: 150 m Speed: 30 kn*10

Sorted: 148 Kg Total catch: 265.85 CATCH/HOUR: 531.70
SPECIES CATCH/HOUR % OF TOT. C SAMP
weight numbers
Chlorophthalmus atlanticus 368.10 10858 34.83 4311
Spicara alta 362.70 1960 34.32 4306
Antigonius capros 162.00 4924 15.33
Dentex angelensis 29.88 224 2.83 4309
Zenopsis conchifer 27.60 20 2.61 4308
Trachurus trecae 24.04 5854 2.27 4307
Dentex congoides 21.34 198 2.02 4304
Todaropsis eblanae 12.42 100 1.18
Squatina oculata 10.00 4 0.95
Zeus faber 8.52 16 0.81 4310
Trichiurus lepturus 5.58 100 0.53
Sphoeroides pacchaster 5.40 8 0.51
Ariomma bondi 3.34 46 0.32
Umbrina canariensis 2.66 8 0.25 4305
Squalus mitsukurii 2.30 2 0.22
Octopus vulgaris 2.00 2 0.19
Lepidotrigla cadmani 1.98 28 0.19
Eceneis naurocrates 1.90 2 0.18
Pterothrius bellucci 1.62 10 0.15
Scorpaena stephanica 1.00 10 0.09
Raja miraletus 0.80 2 0.08
Pentheroscion mbizi 0.72 10 0.07
Synchiropus phaeton 0.64 10 0.06
Serranus africana 0.28 10 0.03
Total 1056.82 100.02

PROJECT STATION: 994
DATE: 1/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 1054
start stop duration Long W 1712
TIME :14:10:31 14:40:58 30 (min) Purpose code: 3
LOG :6763.82 6765.35 1.54 Area code : 1
FDEPTH: 178 152 GearCond.code:
BDEPTH: 178 152 Validity code:
Towing dir: 62° Wire out: 470 m Speed: 30 kn*10
Sorted: 148 Kg Total catch: 265.85 CATCH/HOUR: 531.70
SPECIES CATCH/HOUR % OF TOT. C SAMP
weight numbers
Decapterus rhonchus 158.20 612 29.75 4325
Dasyatis pastinaca 120.00 2 22.57
Pagellus bellottii 109.68 934 20.63 4324
Pseudupeneus prayensis 82.20 770 15.46 4321
Trachurus trecae 31.20 5.87
Epinephelus aeneus 6.16 10 1.16 4322
Seriola sp. 4.20 10 0.79
Pomadasys rogeri 3.40 10 0.64
Acanthurus monroviae 2.40 4 0.45
Plectrohinchus mediterraneus 2.40 4 0.45
Pomadasys jubelini 2.40 4 0.45
Octopus vulgaris 1.80 4 0.34
Priacanthus arenatus 1.64 10 0.31
Sepia officinalis hierredda 1.40 4 0.26
Scomber japonicus 1.02 52 0.19 4323
Mugil sp. 1.00 4 0.19
Pagrus caeruleostictus 0.94 4 0.18
Torpedo torpedo 0.80 4 0.15
Syacium micrurum 0.80 4 0.15
Sardinella aurita 0.14 4 0.03
Total 531.78 100.02

PROJECT STATION: 995
DATE: 1/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 1054
start stop duration Long W 1712
TIME :14:10:31 14:40:58 30 (min) Purpose code: 3
LOG :6763.82 6765.35 1.54 Area code : 1
FDEPTH: 178 152 GearCond.code:
BDEPTH: 178 152 Validity code:
Towing dir: 62° Wire out: 470 m Speed: 30 kn*10
Sorted: 141 Kg Total catch: 528.47 CATCH/HOUR: 1056.94

PROJECT STATION: 998
DATE: 2/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 1042
start stop duration Long W 1629
TIME :11:00:30 11:30:38 30 (min) Purpose code: 3
LOG :6901.60 6903.17 1.56 Area code : 1
FDEPTH: 35 37 GearCond.code:
BDEPTH: 35 37 Validity code:
Towing dir: 245° Wire out: 150 m Speed: 30 kn*10

Sorted: 111 Kg Total catch: 111.51 CATCH/HOUR: 215.83
SPECIES CATCH/HOUR % OF TOT. C SAMP
weight numbers
Chlorophthalmus atlanticus 130.65 3416 43.59 4312
Synagrops microlepis 31.35 1622 10.46
Ilex coindetii 31.32 252 10.45
Zenopsis conchifer 24.39 19 8.14 4314
CHIROSTYLIDAE 22.22 2685 7.41
Pterothrius bellucci 11.54 79 3.85
Todaropsis eblanae 7.41 105 2.47
Hypoclydonia bella 6.54 466 2.18
Ariomma bondi 6.10 139 2.04
Merluccius polli 5.90 31 1.97 4313
Parasudis fraser-bruenneri 4.84 706 1.61
Brotila barbata 3.77 2 1.26
Parapeneus longirostris 3.41 379 1.14
Lepidotrigla cadmani 2.13 39 0.71
Umbrina canariensis 1.43 2 0.48
Trigla lyra 1.34 4 0.45
Raja miraletus 0.97 2 0.32
Pontinus accraensis 0.74 4 0.25
Antigonia capros 0.74 56 0.25

PROJECT STATION: 994
DATE: 1/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 1054
start stop duration Long W 1712
TIME :14:10:31 14:40:58 30 (min) Purpose code: 3
LOG :6763.82 6765.35 1.54 Area code : 1
FDEPTH: 178 152 GearCond.code:
BDEPTH: 178 152 Validity code:
Towing dir: 62° Wire out: 470 m Speed: 30 kn*10
Sorted: 111 Kg Total catch: 111.51 CATCH/HOUR: 215.83
SPECIES CATCH/HOUR % OF TOT. C SAMP
weight numbers
Decapterus rhonchus 127.55 830 59.10 4327
Sardinella aurita 24.00 145 11.12 4326
Fistularia petimba 15.29 7.08
Pagellus bellottii 10.94 89 5.07 4328
Priacanthus arenatus 8.85 72 4.10
Sepia officinalis hierredda 4.66 33 2.16 4331
Decapterus punctatus 3.66 74 1.70 4329
Trichiurus lepturus 3.00 4 1.39
Dactylopterus volitans 2.54 6 1.18
Pseudupeneus prayensis 2.50 19 1.16 4330
Trachinophthalmus myops 2.28 10 1.06
Scomber japonicus 1.84 10 0.85
Trachurus trecae 1.74 15 0.81
Raja miraletus 1.55 4 0.72
Octopus vulgaris 1.39 2 0.64
Lagocephalus laevigatus 1.16 2 0.54
Sphyraena guachancho 0.97 2 0.45
Chelidonichthys gabonensis 0.70 6 0.32
Pagrus caeruleostictus 0.68 2 0.32
Eceneis naurocrates 0.54 2 0.25
Total 215.84 100.02

PROJECT STATION: 1000
DATE: 2/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 1034
start stop duration Long W 1648
TIME :14:43:08 13:14:04 31 (min) Purpose code: 3
LOG :6912.39 6914.00 1.58 Area code : 1
FDEPTH: 59 64 GearCond.code:
BDEPTH: 59 64 Validity code:
Towing dir: 245° Wire out: 200 m Speed: 30 kn*10

Squatina oculata	11.60	2	3.57	SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
Ariomma bondi	7.30	426	2.25	Decapterus punctatus	weight numbers	1059.30	30696
Trichiurus lepturus	5.20	192	1.60	Sardinella aurita	18.81	510	1.72
CHIROSTYLIDAE	5.20	1702	1.60	Scomber japonicus	11.55	180	1.06
Pterothrissus belloci	5.16	46	1.59	Chelidonichthys gabonensis	2.16	18	0.20
Squalus mitsukurii	4.00	4	1.23	Chiroscombrus chrysurus	1.32	51	0.12
Illex coindetii	3.70	70	1.14	Priacanthus arenatus	0.84	18	0.08
Todaropsis ebiana	2.50	100	0.77	Total		1093.98	100.01
Dentex angolensis	1.80	4	0.55				
Parapenaeus longirostris	1.10	206	0.34				
Antigonia capros	0.60	10	0.18				
Heptanchias perlo	0.60	2	0.18				
Aulopus cadenati	0.54	6	0.17				
Merluccius polli	0.30	6	0.09				
Sepia officinalis hierredda	0.20	10	0.06				
Pontinus acraensis	0.18	6	0.06				
Grammoplites gruveli	0.12	6	0.04				
Parasudis fraser-bruenneri	0.10	16	0.03				
Monolete microstoma	0.06	6	0.02				
Total	324.86	100.00					

PROJECT STATION:1013
DATE: 4/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 934
start stop duration Long W 1610

TIME :18:59:41 19:29:48 30 (min) Purpose code: 3
LOG :7273.01 7274.47 1.45 Area code : 2
FDEPTH: 142 133 GearCond.code:
BDEPTH: 142 133 Validity code:
Towing dir: 40° Wire out: 350 m Speed: 30 kn*10

Sorted: 60 Kg Total catch: 211.43 CATCH/HOUR: 422.86

SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
	weight numbers		
Ariomma bondi	177.80	9994	42.05
Antigonia capros	86.10	3826	20.36
Trachurus trecae	77.00		18.21
Illex coindetii	26.04	1066	6.16
Aulopus cadenati	13.24	252	3.13
Syacium micrurum	10.86	162	2.57
Lagocephalus laevigatus	6.72	36	1.59
Uranoscopus albesca	4.80	8	1.14
Raja miraletus	4.08	8	0.96
Liocraninus corrugatus	3.44	722	0.81
Priacanthus arenatus	2.24	42	0.53
Dactylopterus volitans	1.92	8	0.45
Lepidotrigla carolae	1.82	78	0.43
Chelidonichthys gabonensis	1.20	14	0.28
Trichiurus lepturus	1.12	28	0.26
Zeus faber	0.92	36	0.22
Trachinus pellegrini	0.84	14	0.20
Pontinus acraensis	0.70	22	0.17
Grammoplites gruveli	0.64	8	0.15
Lepidotrigla cadiami	0.56	8	0.13
Synchiropus phaeton	0.50	22	0.12
Saurida brasiliensis	0.24	8	0.06
Peristedion cataphractum	0.08	8	0.02
Total	422.86	100.00	

PROJECT STATION:1009
DATE: 4/ 5/06 GEAR TYPE: PT No: 2 POSITION:Lat N 929
start stop duration Long W 1614

TIME :02:25:50 02:56:54 31 (min) Purpose code: 1
LOG :7154.00 7155.69 1.68 Area code : 2
FDEPTH: 40 44 GearCond.code:
BDEPTH: 250 194 Validity code:
Towing dir: 41° Wire out: 100 m Speed: 32 kn*10

Sorted: 22 Kg Total catch: 22.59 CATCH/HOUR: 43.72

SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
	weight numbers		
MYCTOPHIDAE	34.84	79.69	
Shrimps, small, non comm.	3.87	8.85	
Trachipterus trachypterus	3.15	17	7.20
PARALEPIDIDAE	0.79	89	1.81
Ariomma bondi	0.31	19	0.71
Illex coindetii	0.21	6	0.48
Sphoeroides marmoratus	0.14	6	0.32
Trichiurus lepturus	0.14	4	0.32
Hypoclydonia bella	0.14	12	0.32
Synagrops microlepis	0.06	2	0.14
GONOSTOMATIDAE	0.04	4	0.09
Fistularia petimba	0.02	2	0.05
Selene dorsalis, juveniles	0.00	4	
Total	43.71	99.98	

PROJECT STATION:1010
DATE: 4/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 955
start stop duration Long W 1550

TIME :09:19:24 09:49:07 30 (min) Purpose code: 3
LOG :7197.20 7198.81 1.60 Area code : 2
FDEPTH: 36 38 GearCond.code:
BDEPTH: 36 38 Validity code:
Towing dir: 225° Wire out: 150 m Speed: 32 kn*10

Sorted: 148 Kg Total catch: 541.24 CATCH/HOUR: 1082.48

SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
	weight numbers		
Chloroscombrus chrysurus	793.60	16056	73.31
Sardinella maderensis	97.60	1772	9.02
Decapterus punctatus	55.60	2512	5.14
Sardinella aurita	55.28	776	5.11
Decapterus rhonchus	28.08	152	2.59
Sphyraena afra	26.60	2	2.46
Brachydeuterus auritus	9.92	104	0.92
Trachurus trecae	3.68	328	0.34
Balistes punctatus	3.40	4	0.31
Pseudupeneus prayensis	2.76	20	0.25
Pagellus bellottii	2.04	18	0.19
Selene dorsalis	1.12	16	0.10
Pagrus caeruleostictus	1.04	6	0.10
Trachinocephalus myops	0.80	8	0.07
Xyrichtys novacula	0.56	8	0.05
Scomber japonicus	0.40	8	0.04
Total	1082.48	100.00	

PROJECT STATION:1010
DATE: 4/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 1009
start stop duration Long W 1537

TIME :12:23:23 12:53:31 30 (min) Purpose code: 3
LOG :7222.00 7223.47 1.45 Area code : 2
FDEPTH: 26 28 GearCond.code:
BDEPTH: 26 28 Validity code:
Towing dir: 220° Wire out: 150 m Speed: 30 kn*10

Sorted: 108 Kg Total catch: 156.20 CATCH/HOUR: 312.40

SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
	weight numbers		
Chloroscombrus chrysurus	114.40	1952	36.62
Rachycentron canadum	91.40	6	29.26
Decapterus rhonchus	66.60		21.32
Decapterus punctatus	10.60		3.39
Sepia officinalis hierredda	7.20	14	2.30
Sardinella aurita	6.00	362	1.92
Cronius ruber	4.80	88	1.54
Pagrus caeruleostictus	3.12	74	1.00
Decapterus rhonchus	2.88	38	0.92
Sardinella maderensis	2.20	80	0.70
Trachurus trecae	1.80		0.58
Balistes capricrus	1.16	4	0.37
Brachydeuterus auritus	0.16	8	0.05
Pagellus bellottii	0.08	4	0.03
Total	312.40	100.00	

PROJECT STATION:1012
DATE: 4/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 939
start stop duration Long W 1605

TIME :17:40:35 18:00:57 20 (min) Purpose code: 3
LOG :7264.33 7265.53 1.20 Area code : 2
FDEPTH: 68 78 GearCond.code:
BDEPTH: 68 78 Validity code:
Towing dir: 220° Wire out: 220 m Speed: 30 kn*10

Sorted: 66 Kg Total catch: 364.66 CATCH/HOUR: 1093.98

PROJECT STATION:1016
DATE: 5/ 5/06 GEAR TYPE: BT No: 2 POSITION:Lat N 910
start stop duration Long W 1535

TIME :05:12:47 05:43:42 31 (min) Purpose code: 3
LOG :7333.03 7335.00 1.96 Area code : 2
FDEPTH: 40 43 GearCond.code:
BDEPTH: 351 322 Validity code:
Towing dir: 310° Wire out: 130 m Speed: 38 kn*10

Sorted: 17 Kg Total catch: 17.76 CATCH/HOUR: 34.37

SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
	weight numbers		
MYCTOPHIDAE	23.23	8061	67.59
Ariomma bondi	7.35	552	21.38
Trachipterus trachypterus	1.26	14	3.67
Synagrops microlepis	1.16	50	3.38
Hypoclydonia bella	1.01	101	2.94
CARANGIDAE	0.12	48	0.35
PARALEPIDIDAE	0.12	25	0.35
CARANGIDAE	0.06	33	0.17
Selene dorsalis, juveniles	0.06	29	0.17
Illex coindetii	0.02	6	0.06
Total	34.39	100.06	

PROJECT STATION:1016
DATE: 5/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 913
start stop duration Long W 1539

TIME :07:27:34 07:57:47 30 (min) Purpose code: 3

Sardiniella spp. (juv.)	39.00	7242	5.88		Chelidonichthys gabonensis	0.27	4	0.42
Pagrus caeruleostrictus	27.60	2068	4.16	4384	Bothus podas africanus	0.14	6	0.22
Engraulis encrasicolus	24.60	4920	3.71	4385	Trachinus armatus	0.08	2	0.12
Pagellus bellottii	12.20	2604	1.84		Chromis cadehati	0.06	2	0.09
Sphyraena afra	7.40	2	1.11		Total		64.43	100.02
Selar crumenophthalmus	6.74	80	1.02					
Decapterus rhonchus	5.80	30	0.87					
Sepla officinalis hierredda	3.60	8	0.54					
Ephippion guttifer	3.58	2	0.54					
Sphyraena guachancho	2.86	8	0.43					
Sepia officinalis hierredda	2.30	76	0.35					
Chloroscombrus chrysurus	2.20	36	0.33					
Lagocephalus laevigatus	2.06	6	0.31					
Fistularia petimba	2.04	36	0.31					
Chaetodipterus gorenensis	2.00	8	0.30					
Psettodes belcheri	1.92	6	0.29					
Portunus validus	1.78	4	0.27					
Rachycentron canadum	1.70	2	0.26					
Sphyraena guachancho	1.54	44	0.23					
Caranx cryos	0.90	8	0.14					
Selene dorsalis	0.90	14	0.14					
Pseudupeneus prayensis	0.80	154	0.12					
Chloroscombrus chrysurus	0.80	306	0.12					
Sardiniella maderensis	0.58	14	0.09					
Priacanthus arenatus	0.48	8	0.07					
Total	663.78	100.02						

PROJECT STATION:1025								
DATE: 6/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 933								
start stop duration								
TIME :09:25:39 09:45:38 20 (min) Purpose code: 3								
LOG :7504.00 7505.08 1.07 Area code : 2								
FDEPTH: 35 32 GearCond.code:								
BDEPTH: 35 32 Validity code:								
Towing dir: 220° Wire out: 150 m Speed: 30 kn*10								
Sorted: 39 Kg	Total catch:	603.83	CATCH/HOUR:	1811.49				

PROJECT STATION:1025								
DATE: 6/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 933								
start stop duration								
TIME :09:25:39 09:45:38 20 (min) Purpose code: 3								
LOG :7504.00 7505.08 1.07 Area code : 2								
FDEPTH: 35 32 GearCond.code:								
BDEPTH: 35 32 Validity code:								
Towing dir: 220° Wire out: 150 m Speed: 30 kn*10								
Sorted: 39 Kg	Total catch:	603.83	CATCH/HOUR:	1811.49				

PROJECT STATION:1026								
DATE: 6/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 916								
start stop duration								
TIME :13:00:36 13:30:23 30 (min) Purpose code: 3								
LOG :7529.33 7530.89 1.55 Area code : 2								
FDEPTH: 48 40 GearCond.code:								
BDEPTH: 48 40 Validity code:								
Towing dir: 40° Wire out: 150 m Speed: 30 kn*10								
Sorted: 320 Kg	Total catch:	320.25	CATCH/HOUR:	640.50				

PROJECT STATION:1026								
DATE: 6/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 916								
start stop duration								
TIME :13:00:36 13:30:23 30 (min) Purpose code: 3								
LOG :7529.33 7530.89 1.55 Area code : 2								
FDEPTH: 48 40 GearCond.code:								
BDEPTH: 48 40 Validity code:								
Towing dir: 40° Wire out: 150 m Speed: 30 kn*10								
Sorted: 320 Kg	Total catch:	320.25	CATCH/HOUR:	640.50				

PROJECT STATION:1027								
DATE: 6/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 904								
start stop duration								
TIME :15:59:35 16:30:59 31 (min) Purpose code: 3								
LOG :7550.83 7552.42 1.57 Area code : 3								
FDEPTH: 73 66 GearCond.code:								
BDEPTH: 73 66 Validity code:								
Towing dir: 40° Wire out: 190 m Speed: 30 kn*10								
Sorted: 33 Kg	Total catch:	33.29	CATCH/HOUR:	64.43				

PROJECT STATION:1027								
DATE: 6/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 882								
start stop duration								
TIME :02:18:52 02:49:06 30 (min) Purpose code: 1								
LOG :7620.19 7621.75 1.55 Area code : 3								
FDEPTH: 15 35 GearCond.code:								
BDEPTH: 107 93 Validity code:								
Towing dir: 40° Wire out: 130 m Speed: 30 kn*10								
Sorted: 17 Kg	Total catch:	17.17	CATCH/HOUR:	34.34				

PROJECT STATION:1030								
DATE: 7/ 5/06 GEAR TYPE: PT No: 7 POSITION:Lat N 832								
start stop duration								
TIME :03:54:35 04:25:07 31 (min) Purpose code: 1								
LOG :7627.69 7629.40 1.70 Area code : 3								
FDEPTH: 35 39 GearCond.code:								
BDEPTH: 66 79 Validity code:								
Towing dir: 220° Wire out: 130 m Speed: 30 kn*10								
Sorted: 5 Kg	Total catch:	5.51	CATCH/HOUR:	10.66				

PROJECT STATION:1031								
DATE: 7/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 832								
start stop duration								
TIME :07:03:06 07:32:58 30 (min) Purpose code: 3</								

PROJECT STATION:1032
DATE: 7/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 834
start stop duration Long W 1420
TIME :10:33:33 11:03:36 30 (min) Purpose code: 3
LOG :7658.51 7660.15 1.63 Area code : 3
FDEPTH: 48 59 GearCond.code:
BDEPTH: 48 59 Validity code:
Towing dir: 225° Wire out: 180 m Speed: 30 kn*10
Sorted: 46 Kg Total catch: 444.52 CATCH/HOUR: 889.04

SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
	weight numbers		
Decapterus rhonchus	482.40	3334	54.26
Chloroscombrus chrysurus	266.40	3618	29.96
Decapterus punctatus	96.72	1364	10.88
Sardinella maderensis	13.92	98	1.57
Lagocephalus laeavigatus	6.24	72	0.70
Dactylopterus volitans	5.74	10	0.65
Priacanthus arenatus	5.22	98	0.59
Pseudupeneus prayensis	2.70	36	0.30
Selene dorsalis	2.40	24	0.27
Raja miraletus	1.62	8	0.18
Bothus podas africanus	1.20	24	0.13
Fistularia petimba	0.98	18	0.11
Sepia officinalis hierredda	0.94	6	0.11
Caranx cryos	0.94	4	0.11
Torpedo torpedo	0.64	2	0.07
Pagrus caeruleostictus	0.60	2	0.07
Xyrichtys novacula	0.38	10	0.04
Total	889.04	100.00	

PROJECT STATION:1036
DATE: 7/ 5/06 GEAR TYPE: PT No: 7 POSITION:Lat N 911
start stop duration Long W 1349
TIME :22:37:04 23:08:03 31 (min) Purpose code: 1
LOG :7739.62 7741.41 1.79 Area code : 3
FDEPTH: 20 15 GearCond.code:
BDEPTH: 30 25 Validity code:
Towing dir: 175° Wire out: 130 m Speed: 30 kn*10
Sorted: 45 Kg Total catch: 215.07 CATCH/HOUR: 416.26

SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
	weight numbers		
Sardinella aurita	347.23	27573	83.42
Decapterus punctatus	33.45	2843	8.04
Sphyraena afra	20.13	2	4.84
Sardinella maderensis	11.50	105	2.76
Scomberomorus tritor	2.11	4	0.51
Sepia officinalis hierredda	0.70	12	0.17
Decapterus rhonchus	0.70	70	0.17
Fistularia petimba	0.35	12	0.08
Engraulis encrasicolus	0.12	46	0.03
Total	416.29	100.02	

PROJECT STATION:1033
DATE: 7/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 849
start stop duration Long W 1408
TIME :15:16:19 15:46:02 30 (min) Purpose code: 3
LOG :7692.20 7693.75 1.55 Area code : 3
FDEPTH: 38 40 GearCond.code:
BDEPTH: 38 40 Validity code:
Towing dir: 220° Wire out: 150 m Speed: 30 kn*10
Sorted: 118 Kg Total catch: 118.86 CATCH/HOUR: 237.72

SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
	weight numbers		
Pagrus caeruleostictus	75.40	512	31.72
Pagellus bellottii	60.00	378	25.24
Pseudupeneus prayensis	42.00	550	17.67
Acanthurus monroviae	13.00	10	5.47
Chloroscombrus chrysurus	11.24	148	4.73
Bodianus speciosus	5.68	2	2.39
Dactylopterus volitans	5.44	6	2.29
Aluterus heudelotii	5.20	12	2.19
Caranx cryos	3.80	6	1.60
Diodon holocanthus	3.20	6	1.35
Fistularia petimba	3.14	100	1.32
Raja miraletus	3.00	8	1.26
Sepia officinalis hierredda	2.78		1.17
Sphyraena sphyraena	1.20	4	0.50
Trachinocephalus myops	0.94	4	0.40
Xyrichtys novacula	0.42	6	0.18
Chilomycterus spinosus mauret.	0.42	2	0.18
Lagocephalus laeavigatus	0.34	2	0.14
Chaetodon hoefleri	0.26	2	0.11
Sparusoma rubripinne	0.26	2	0.11
Total	237.72	100.02	

PROJECT STATION:1034
DATE: 7/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 902
start stop duration Long W 1358
TIME :18:02:35 18:33:04 30 (min) Purpose code: 3
LOG :7713.76 7715.58 1.81 Area code : 3
FDEPTH: 28 32 GearCond.code:
BDEPTH: 28 32 Validity code:
Towing dir: 220° Wire out: 150 m Speed: 30 kn*10
Sorted: 226 Kg Total catch: 276.94 CATCH/HOUR: 553.88

SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
	weight numbers		
Pomadasys rogeri	241.60	276	43.62
Pagrus caeruleostictus	99.40	600	17.95
Sardinella aurita	71.60	4406	12.93
Alectis alexandrinus	44.00	44	7.94
Decapterus punctatus	39.20	2864	7.08
Pagellus bellottii	30.00	256	5.42
Acanthurus monroviae	5.68	6	1.03
Bodianus speciosus	5.60	4	1.01
Epinephelus aeneus	4.20	6	0.76
Aluterus monoceros	2.80	6	0.51
Balistes punctatus	2.20	2	0.40
Lutjanus goreensis	1.80	2	0.32
Sepia officinalis hierredda	1.50	4	0.27
Lethrinus atlanticus	1.40	2	0.25
Chaetodipterus goreensis	1.12	2	0.20
Torpedo torpedo	1.06	2	0.19
Lutjanus fulgens	0.60	2	0.11
Pseudupeneus prayensis	0.12	10	0.02
Total	553.88	100.01	

PROJECT STATION:1035
DATE: 7/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 914
start stop duration Long W 1348
TIME :20:51:49 21:23:24 32 (min) Purpose code: 3
LOG :7735.49 7737.27 1.76 Area code : 3
FDEPTH: 25 26 GearCond.code:
BDEPTH: 25 26 Validity code:
Towing dir: 220° Wire out: 130 m Speed: 30 kn*10
Sorted: 94 Kg Total catch: 115.46 CATCH/HOUR: 216.49

SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
	weight numbers		
Pagrus caeruleostictus	71.44	842	33.00
Psettosideres sp.	34.88	139	16.11
Ephippion guttifer	26.63	11	12.30
Pagellus bellottii	18.56	152	8.57
Pseudupeneus prayensis	15.41	1453	7.12
Decapterus punctatus	12.71	671	5.87
Decapterus rhonchus	11.03	1592	5.09
Decapterus rhonchus	5.81	17	2.68
Chaetodipterus goreensis	3.75	11	1.73
Penaeus notialis	3.06	86	1.41
Total	349.98	100.00	

PROJECT STATION:1039
DATE: 8/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 830
start stop duration Long W 1347
TIME :07:24:03 07:54:01 30 (min) Purpose code: 3
LOG :7789.07 7790.93 1.84 Area code : 3
FDEPTH: 31 33 GearCond.code:
BDEPTH: 31 33 Validity code:
Towing dir: 215° Wire out: 150 m Speed: 30 kn*10
Sorted: 164 Kg Total catch: 486.34 CATCH/HOUR: 972.68

SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
	weight numbers		
Priacanthus arenatus	2.81	68	1.30
Brachydeuterus auritus	1.80	225	0.83
Conger conger	1.54	4	0.71
Sepia officinalis hierredda	1.31	8	0.61
Syacium micrurum	1.13	23	0.52
Scorpaena scrofa	1.03	2	0.48
Penaeus kerathurus	0.75	36	0.35
Cronius ruber	0.68	11	0.31
Sardinella aurita	0.68	68	0.31
Selar crumenophthalmus	0.47	8	0.22
Raja miraletus	0.38	2	0.18
Torpedo torpedo	0.32	2	0.15
Grammoplites griseus	0.11	11	0.05
Engraulis encrasicolus	0.11	23	0.05
Trachinocephalus myops	0.11	2	0.05
Total	216.51	100.00	

	weight	numbers	Total	1221.96	100.01
Decapterus punctatus	531.00	50066	54.59	4451	
Sardinella aurita	116.00	9592	11.93	4448	
Pagellus bellottii	106.20	884	10.92	4447	
Pagrus caeruleostictus	63.80	238	6.56	4449	
Lethrinus atlanticus	38.80	72	3.99	4450	
Acanthurus monroviae	30.20	50	3.10		
Cymbium sp.	24.00	6	2.47		
Sepia officinalis hierredda	14.94	22	1.54	4452	
Balistes punctatus	12.70	14	1.31		
Chaetodipterus goreensis	8.20		0.84		
Pseudupeneus prayensis	7.20	70	0.74	4446	
Brachydeuterus auritus	7.00	100	0.72		
Bodianus speciosus	3.36	6	0.35		
Decapterus rhonchus	3.00	100	0.31		
Caranx cryos	3.00	2	0.31		
Psettodes belcheri	1.82	4	0.19		
Epinephelus costae	1.46	2	0.15		
Total	972.68	100.02			

PROJECT STATION:1043
DATE: 8/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 809
start stop duration Long W 1402
TIME :13:37:41 14:08:01 30 (min) Purpose code: 3
LOG :7821.14 7822.64 1.49 Area code : 3
FDEPTH: 26 26 GearCond.code:
BDEPTH: 26 26 Validity code:
Towing dir: 215° Wire out: 150 m Speed: 30 kn*10

Sorted: 123 Kg Total catch: 123.98 CATCH/HOUR: 247.96

	weight	numbers	SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
Balistes capriscus	90.80	100		36.62	4475	
Chloroscombrus chrysurus	80.80	1692		32.59	4479	
Dactylopterus volitans	47.60	166		19.20		
Decapterus rhonchus	10.44	60		4.21	4478	
Pagrus caeruleostictus	9.08	72		3.66	4477	
Pseudupeneus prayensis	2.24	16		0.90	4480	
Diodon holocanthus	1.64	4		0.66		
Chilomycterus spinosus mauret.	0.90	4		0.36		
Sepia officinalis hierredda	0.74	2		0.30		
Aluterus monoceros	0.68	2		0.27		
Pagellus bellottii	0.64	4		0.26		
Priacanthus arenatus	0.58	8		0.23		
Eucinostomus melanopterus	0.52	6		0.21		
Selar crumenophthalmus	0.46	10		0.19		
Echeneis naucrates	0.38	2		0.15		
Trachinocephalus myops	0.20	2		0.08		
Xyrichtys novacula	0.18	2		0.07		
Sardinella maderensis	0.08	2		0.03		
Total	972.68	100.02				

	weight	numbers	SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
Chloroscombrus chrysurus	1250.00	36658		69.65	4458	
Brachydeuterus auritus	255.20	8210		14.22	4459	
Decapterus rhonchus	106.60	436		5.94	4453	
Pagellus bellottii	68.40	952		3.81	4461	
Decapterus rhonchus	21.40	666		1.19	4457	
Caranx cryos	17.40	42		0.97	4455	
Selene dorsalis	13.60	290		0.76		
Balistes capriscus	11.80	8		0.66		
Sphyraena guachancho	7.80	58		0.43		
Selar crumenophthalmus	6.30	24		0.35	4454	
Pseudupeneus prayensis	5.74	76		0.32	4460	
Boops boops	4.94	88		0.28		
Pagrus caeruleostictus	4.66	44		0.26	4456	
Decapterus punctatus	4.20	88		0.23		
Dactylopterus volitans	4.20	6		0.23		
Sardinella maderensis	3.20	30		0.18		
Euthynnus alletteratus	2.32	4		0.13		
Priacanthus arenatus	2.00	58		0.11		
Caranx cryos	1.80	30		0.10		
Sepia officinalis hierredda	1.74	2		0.10		
Sphyraena guachancho	0.58	4		0.03		
Sphyraena sphyraena	0.50	2		0.03		
Sphyraena guachancho	0.48	10		0.03		
Galeoides decadactylus	0.32	2		0.02		
Total	1795.18	100.03				

	weight	numbers	SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
Chloroscombrus chrysurus	1250.00	36658		69.65	4458	
Brachydeuterus auritus	255.20	8210		14.22	4459	
Decapterus rhonchus	106.60	436		5.94	4453	
Pagellus bellottii	68.40	952		3.81	4461	
Decapterus rhonchus	21.40	666		1.19	4457	
Caranx cryos	17.40	42		0.97	4455	
Selene dorsalis	13.60	290		0.76		
Balistes capriscus	11.80	8		0.66		
Sphyraena guachancho	7.80	58		0.43		
Selar crumenophthalmus	6.30	24		0.35	4454	
Pseudupeneus prayensis	5.74	76		0.32	4460	
Boops boops	4.94	88		0.28		
Pagrus caeruleostictus	4.66	44		0.26	4456	
Decapterus punctatus	4.20	88		0.23		
Dactylopterus volitans	4.20	6		0.23		
Sardinella maderensis	3.20	30		0.18		
Euthynnus alletteratus	2.32	4		0.13		
Priacanthus arenatus	2.00	58		0.11		
Caranx cryos	1.80	30		0.10		
Sepia officinalis hierredda	1.74	2		0.10		
Sphyraena guachancho	0.58	4		0.03		
Sphyraena sphyraena	0.50	2		0.03		
Sphyraena guachancho	0.48	10		0.03		
Galeoides decadactylus	0.32	2		0.02		
Total	1795.18	100.03				

	weight	numbers	SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
Chloroscombrus chrysurus	2690.00	45154		88.26	4483	
Pagellus bellottii	146.60	1040		4.81	4487	
Pseudupeneus prayensis	74.40	680		2.44	4482	
Pagrus caeruleostictus	29.60	180		0.97	4485	
Selene dorsalis	20.00	380		0.66	4484	
Chaetodipterus goreensis	13.20	32		0.43	4481	
Pomadasys rogeri	12.30	22		0.40	4486	
Caranx cryos	11.40	20		0.37		
Selar crumenophthalmus	11.20	60		0.37		
Dactylopterus volitans	10.80	40		0.35		
Lagocephalus laevigatus	9.40	60		0.31		
Decapterus rhonchus	6.80	20		0.22		
Eucinostomus melanopterus	3.20	40		0.10		
Brachydeuterus auritus	2.80	20		0.09		
Aluterus heudelotii	1.80	20		0.06		
Priacanthus arenatus	1.80	20		0.06		
Sphyraena guachancho	1.52	4		0.05		
Ephippion guttifer	1.10	2		0.04		
Total	3047.92	99.99				

	weight	numbers	SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
Chloroscombrus chrysurus	2690.00	45154		88.26	4483	
Pagellus bellottii	146.60	1040		4.81	4487	
Pseudupeneus prayensis	74.40	680		2.44	4482	
Pagrus caeruleostictus	29.60	180		0.97	4485	
Selene dorsalis	20.00	380		0.66	4484	
Chaetodipterus goreensis	13.20	32		0.43	4481	
Pomadasys rogeri	12.30	22		0.40	4486	
Caranx sennegallus	8.00	8		0.37		
Pagellus bellottii	5.80	54		1.10	4491	
Sepia officinalis hierredda	3.00	6		0.57		
Brachydeuterus auritus	1.84	20		0.35	4493	
Selar crumenophthalmus	1.80	6		0.34		
Decapterus rhonchus	1.12	6		0.21		
Selene dorsalis	0.92	46		0.17	4494	
Aluterus heudelotii	0.48	6		0.09		
Lagocephalus laevigatus	0.24	2		0.05		
Total	3047.92	99.99				

	weight	numbers	SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
Rhizoprionodon acutus	213.20	122		40.46		
Dactylopterus volitans	133.60	506		25.36		
Balistes capriscus	103.60	92		19.66	4489	
Chloroscombrus chrysurus	21.60	412		4.10	4492	
Chaetodipterus goreensis	16.70	38		3.17	4488	
Pseudupeneus prayensis	10.00	104		1.90	4490	
Caranx sennegallus	8.00	8		1.52		
Pagellus bellottii	5.80	54		1.10	4491	
Pomadasys rogeri	4.80	8		0.91		
Sepia officinalis hierredda	3.00	6		0.57		
Brachydeuterus auritus	1.84	20		0.35	4493	
Selar crumenophthalmus	1.80	6		0.34		
Decapterus rhonchus	1.12	6		0.21		
Selene dorsalis	0.92	46		0.17	4494	
Aluterus heudelotii	0.48	6		0.09		
Lagocephalus laevigatus	0.24	2		0.05		
Total	526.70	99.96				

	weight	numbers	SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
DATE: 8/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 814						
start stop duration						
TIME :12:38:45 12:38:50 30 (min) Purpose code: 3						
LOG :7814.70 7816.19 1.79 Area code : 3						
FDEPTH: 25 24 GearCond.code:						
BDEPTH: 25 24 Validity code:						
Towing dir: 215° Wire out: 150 m Speed: 30 kn*10						
Sorted: 165 Kg Total catch: 611.00 CATCH/HOUR: 1222.00						
SPECIES	CATCH/HOUR	% OF TOT. C	SAMP			
Pagrus caeruleostictus	499.20	3404	40.85	4474		
Pagellus bellottii	284.00	1840	23.24	4468		
Chloroscombrus chrysurus	152.00	3578	12.44	4473		
Priacanthus arenatus	129.60					

Pagrus caeruleostictus 9.73 28 0.39 4497
 Dactylopterus volitans 6.44 19 0.26
 Pseudupeneus prayensis 2.78 41 0.11
 Boops boops 2.78 41 0.11
 Dentex canariensis 2.05 3 0.08
 Scorpaena angolensis 1.23 6 0.05
 Dentex angolensis 1.04 6 0.04
 Total 2502.26 100.00

PROJECT STATION:1051
 DATE: 9/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 731
 start stop duration Long W 1341
 TIME :11:05:03 11:35:07 30 (min) Purpose code: 3
 LOG :7935.29 7936.82 1.52 Area code : 3
 FDEPTH: 81 75 GearCond.code:
 BDEPTH: 81 75 Validity code:
 Towing dir: 35ø Wire out: 220 m Speed: 30 kn*10
 Sorted: 289 Kg Total catch: 289.53 CATCH/HOUR: 579.06

PROJECT STATION:1047
 DATE: 9/ 5/06 GEAR TYPE: PT No: 1 POSITION:Lat N 727
 start stop duration Long W 1344
 TIME :01:20:38 01:55:45 35 (min) Purpose code: 1
 LOG :7895.87 7897.98 2.10 Area code : 3
 FDEPTH: 22 29 GearCond.code:
 BDEPTH: 152 89 Validity code:
 Towing dir: 35ø Wire out: 130 m Speed: 30 kn*10
 Sorted: 52 Kg Total catch: 52.50 CATCH/HOUR: 90.00

SPECIES CATCH/HOUR % OF TOT. C SAMP
 weight numbers
 Hypoclydonia bella 48.09 4181 53.43
 Euthynnus alletteratus 27.26 26 30.29 4501
 Alectis alexandrinus 4.37 2 4.86
 Ariomma bondi 3.79 147 4.21 4502
 Echeneis naucrates 3.31 3 3.68
 Illex coindetii 1.47 86 1.63
 Caranx hippos 1.41 2 1.57
 Promethichthys prometheus 0.15 3 0.17
 Saurida brasiliensis 0.10 17 0.11
 Caranx cryos 0.05 2 0.06
 Total 90.00 100.01

SPECIES CATCH/HOUR % OF TOT. C SAMP
 weight numbers
 Decapterus punctatus 240.00 13434 41.45 4508
 Dentex congensis 85.60 996 14.78 4506
 Trachurus trecae 82.00 444 14.16 4511
 Chromis ctenocephala 61.20 538 10.57
 Dentex angolensis 32.40 230 5.60 4509
 Epinephelus aeneus 21.60 2 3.73 4512
 Scomber japonicus 19.40 3.35
 Priacanthus arenatus 11.60 16 2.00
 Pagrus caeruleostictus 9.60 18 1.66 4510
 Boops boops 6.82 90 1.18
 Fistularia petimba 1.94 12 0.34
 Erythrocles monodi 1.20 164 0.21
 Echeneis naucrates 1.20 4 0.21
 Sardinella aurita 0.94 52 0.16 4507
 Decapterus rhonchus 0.86 2 0.15
 Scorpaena sp. 0.74 4 0.13
 Pagellus bellottii 0.74 6 0.13
 Raja miraletus 0.62 2 0.11
 Pseudupeneus prayensis 0.40 4 0.07
 Citharus linguatula 0.20 6 0.03
 Total 579.06 100.02

PROJECT STATION:1048
 DATE: 9/ 5/06 GEAR TYPE: PT No: 1 POSITION:Lat N 740
 start stop duration Long W 1335
 TIME :03:47:08 04:17:05 30 (min) Purpose code: 1
 LOG :7911.60 7913.44 1.82 Area code : 3
 FDEPTH: 47 12 GearCond.code:
 BDEPTH: 47 31 Validity code:
 Towing dir: 35ø Wire out: 130 m Speed: 30 kn*10
 Sorted: 31 Kg Total catch: 35.82 CATCH/HOUR: 71.64

SPECIES CATCH/HOUR % OF TOT. C SAMP
 weight numbers
 Sardinella maderensis 54.80 712 76.49 4503
 Brachydeuterus auritus 7.44 410 10.39 4505
 Chloroscombrus chrysurus 3.98 82 5.56 4504
 Euthynnus alletteratus 3.56 6 4.97
 Decapterus rhonchus 0.88 10 1.23
 Sphyraena guachancho 0.48 10 0.67
 Selene dorsalis 0.40 10 0.56
 Echeneis naucrates 0.10 2 0.14
 Total 71.64 100.01

PROJECT STATION:1052
 DATE: 9/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 716
 start stop duration Long W 1257
 TIME :17:24:40 17:50:41 26 (min) Purpose code: 3
 LOG :7992.35 7993.57 1.06 Area code : 3
 FDEPTH: 73 79 GearCond.code:
 BDEPTH: 73 79 Validity code:
 Towing dir: 225ø Wire out: 200 m Speed: 30 kn*10
 Sorted: 182 Kg Total catch: 182.23 CATCH/HOUR: 420.53

PROJECT STATION:1049
 DATE: 9/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 742
 start stop duration Long W 1333
 TIME :07:28:19 07:58:02 30 (min) Purpose code: 3
 LOG :7918.89 7920.57 1.66 Area code : 3
 FDEPTH: 23 32 GearCond.code:
 BDEPTH: 23 32 Validity code:
 Towing dir: 215ø Wire out: 140 m Speed: 30 kn*10
 Sorted: 163 Kg Total catch: 315.82 CATCH/HOUR: 631.64

SPECIES CATCH/HOUR % OF TOT. C SAMP
 weight numbers
 Chloroscombrus chrysurus 201.60 4814 31.92 4515
 Brachydeuterus auritus 122.60 6248 19.41 4514
 Galeoides decadactylus 104.00 1810 16.47 4516
 Sphyraena guachancho 31.30 68 4.96 4510
 Galeoides decadactylus 28.30 74 4.48 4509
 Arius heudeleti 26.30 10 4.16
 Pagrus caeruleostictus 21.90 140 3.47 4507
 Selene dorsalis 17.70 506 2.80
 Sphyraena guachancho 11.14 170 1.76 4511
 Albula vulpes 8.56 14 1.36 4508
 Rhizoprionodon acutus 8.20 12 1.30
 Pagrus caeruleostictus 6.40 1.01
 Pseudotolithus senegalensis 5.70 6 0.90
 Pomadasys jubellini 5.66 14 0.90 4506
 Eucinostomus melanopterus 5.06 80 0.80
 Sardineella maderensis 4.06 80 0.64 4512
 Pseudupeneus prayensis 3.74 64 0.59 4513
 Total 631.64 100.02

SPECIES CATCH/HOUR % OF TOT. C SAMP
 weight numbers
 Dentex congensis 322.85 3192 76.77 4520
 Pagrus caeruleostictus 24.46 60 5.82 4517
 Dentex angolensis 15.58 99 3.70 4515
 Priacanthus arenatus 11.19 125 2.66
 Squatina oculata 10.78 7 2.56
 Pagellus bellottii 8.54 28 2.03 4516
 Umbrina canariensis 5.54 16 1.32 4518
 Dentex canariensis 4.62 5 1.10 4514
 Trachurus trecae 3.55 25 0.84
 Pseudupeneus prayensis 2.49 42 0.59 4519
 Dentex gibbosus 2.31 2 0.55 4513
 Scorpaena angolensis 1.48 5 0.35
 Zeus faber 1.18 5 0.28
 Sepia officinalis hierredda 1.15 2 0.27
 Ariomma bondi 1.15 53 0.27
 Lutjanus fulgens 0.97 2 0.23
 Raja miraletus 0.72 5 0.17
 Anthias anthias 0.65 83 0.15
 Lagocephalus laevigatus 0.60 5 0.14
 Fistularia petimba 0.42 2 0.10
 Alioteuthis africana 0.12 62 0.03
 Chaetodon robustus 0.12 2 0.03
 Scyllarides herklotsii 0.09 2 0.02
 Lepidotrigla cadmanii 0.09 2 0.02
 Calappa pelii 0.05 2 0.01
 Lepidotrigla carolae 0.02 2
 Total 420.72 100.01

PROJECT STATION:1050
 DATE: 9/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 737
 start stop duration Long W 1337
 TIME :09:29:07 09:44:00 15 (min) Purpose code: 3
 LOG :7925.80 7926.60 0.78 Area code : 3
 FDEPTH: 57 60 GearCond.code: 4
 BDEPTH: 57 60 Validity code: 9
 Towing dir: 215ø Wire out: 180 m Speed: 30 kn*10
 Sorted: Kg Total catch: CATCH/HOUR:

SPECIES CATCH/HOUR % OF TOT. C SAMP
 weight numbers
 J E L L Y F I S H 0.00

PROJECT STATION:1053
 DATE: 9/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 723
 start stop duration Long W 1253
 TIME :18:55:50 19:25:57 30 (min) Purpose code: 3
 LOG :8002.34 8004.00 1.67 Area code : 3
 FDEPTH: 49 43 GearCond.code:
 BDEPTH: 49 43 Validity code:
 Towing dir: 35ø Wire out: 150 m Speed: 30 kn*10
 Sorted: 150 Kg Total catch: 471.99 CATCH/HOUR: 943.98

SPECIES CATCH/HOUR % OF TOT. C SAMP
 weight numbers
 Brachydeuterus auritus 231.60 9608 24.53 4525
 Sphyraena guachancho 180.20 532 19.09 4521
 Pteroscion peli 125.60 4186 13.31
 Ilisha africana 106.80 3132 11.31
 Selene dorsalis 98.40 6560 10.42 4524
 Trichirurus lepturus 62.00 2018 6.57
 Ethmalosa fimbriata 28.56 4106 3.03
 Chloroscombrus chrysurus 20.16 2.14
 Pseudotolithus senegalensis 19.80 46 2.10 4522
 Sphyraena afra 7.30 2 0.77
 Penaeus notialis 7.20 228 0.76 4527
 Arius heudeleti 6.80 4 0.72
 Pentanemus quinquearius 6.48 96 0.69
 Scomberomorus tritor 6.18 10 0.65 4523
 Elops senegalensis 5.76 24 0.61
 Galeoides decadactylus 5.52 240 0.58 4526
 Portunus validus 5.20 24 0.55
 Drepang africana 3.80 2 0.40
 Caranx senegallus 3.10 4 0.33
 Trachurus trecae 2.88 24 0.31
 Selar crumenophthalmus 2.40 24 0.25
 Eucinostomus melanopterus 1.92 48 0.20
 Galeoides decadactylus 1.86 8 0.20
 Serranus accraensis 0.96 48 0.10
 Dicologoglossa cuneata 0.96 24 0.10
 Shrimps, small, non comm. 0.96 408 0.10
 Pseudotolithus senegalensis 0.72 48 0.08
 Pseudotolithus elongatus 0.72 2 0.08
 Penaeus kerathurus 0.14 12 0.01 4528
 Total 943.98 99.99

PROJECT STATION:1054
 DATE: 9/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 728
 start stop duration Long W 1249

TIME :20:42:37 21:12:51 30 (min) Purpose code: 3
LOG :8011.02 8012.59 1.57 Area code : 3
FDEPTH: 19 24 GearCond.code:
BDEPTH: 19 24 Validity code:
Towing dir: 215° Wire out: 140 m Speed: 30 kn*10

Sorted: 95 Kg Total catch: 244.67 CATCH/HOUR: 489.34

SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
	weight	numbers	
Galeoides decadactylus	77.44	3248	15.83 4532
Brachydeuterus auritus	73.44	2476	15.01 4536
Pentanemus quinquareius	55.04	1074	11.25 4537
Pteroscion peli	34.24	1388	7.00
Polydactylus quadrifilis	28.80	2	5.89
Selene dorsalis	24.48	2938	5.00
Pomadasys jubelini	24.40	20	4.99 4533
Galeoides decadactylus	22.80	90	4.66 4534
Lutjanus dentatus	22.40	2	4.58
Ilisha africana	16.32		3.34
Trichiurus lepturus	14.40	304	2.94
Pseudotolithus senegalensis	13.60	34	2.78 4531
Dasyatis margarita	12.80	16	2.62
Chloroscombrus chrysurus	10.88	1088	2.22
Pseudotolithus senegalensis	8.48	112	1.73
Sphyraena guachancho	6.80	16	1.39
Pseudotolithus typus	4.80	2	0.98
Parapenaeopsis atlantica	4.40	1274	0.90
Eucinostomus melanopterus	4.32	64	0.88
Pseudotolithus elongatus	4.20	20	0.86 4535
Scomberomorus tritor	3.54	4	0.72
Arius heudeleti	3.10	2	0.63
Penaeus kerathurus	2.14	96	0.44 4530
Cynoglossus senegalensis	2.08	16	0.43
Drepane africana	1.92	16	0.39
Trachinocephalus myops	1.92	32	0.39
Lethrinus atlanticus	1.80	4	0.37
Drepane africana	1.80	2	0.37
Pseudupeneus prayensis	1.60	16	0.33
Pseudotolithus epipericus	1.60	16	0.33
Caranx senegallus	1.32	2	0.27
Etmalosa fimbriata	0.90	8	0.18
Rhizoprionodon acutus	0.64	2	0.13
Pomadasys peroteti	0.48	2	0.10
Portunus validus	0.34	2	0.07
Penaeus notialis	0.12	2	0.02 4529
Total	489.34	100.02	

Galeoides decadactylus	19.00	4.77
Pteroscion peli	17.50	4.39
Brachydeuterus auritus	17.20	4.32 4551
Caranx crysos	17.10	4.29
Galeoides decadactylus	12.20	70 3.06 4545
Sardinella maderensis	11.30	1424 2.84 4549
Pentanemus quinquareius	11.30	620 2.84 4547
Sphyraena guachancho	10.80	34 2.71 4546
Panulirus regius	6.56	14 1.65
Selene dorsalis	4.90	400 1.23
Caranx crysos	4.00	20 1.00
Sphyraena guachancho	3.20	20 0.80
Drepane africana	2.58	2 0.65
Penaeus notialis	2.42	34 0.61 4553
Cynoglossus senegalensis	2.30	10 0.58
Pseudotolithus moorii	1.90	10 0.48
Portunus validus	1.34	4 0.34
Drepane africana	1.20	50 0.30
Penaeus kerathurus	0.70	48 0.18 4552
Sardinella maderensis	0.62	6 0.16
Etmalosa fimbriata	0.50	2 0.13
Pomadasys jubelini	0.40	10 0.10
Pseudotolithus elongatus	0.40	2 0.10
Pseudotolithus typus	0.32	2 0.08
Microchirus wittei	0.20	10 0.05
Total	398.44	100.03

PROJECT STATION:1056
DATE:10/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 713
start stop duration Long W 1219
TIME :08:14:13 08:43:38 29 (min) Purpose code: 3
LOG :8072.77 8074.36 1.58 Area code : 3
FDEPTH: 38 46 GearCond.code:
BDEPTH: 38 46 Validity code:
Towing dir: 215° Wire out: 150 m Speed: 30 kn*10

Sorted: 91 Kg Total catch: 258.75 CATCH/HOUR: 535.34

SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
	weight	numbers	
Brachydeuterus auritus	202.76	6329	37.87
Ilisha africana	61.99	1922	11.58
Sphyraena guachancho	46.24	273	8.64 4558
Sphyraena guachancho	38.81	1260	7.25 4559
Galeoides decadactylus	33.46		6.25
Sardinella maderensis	26.65	1767	4.98 4555
Pseudotolithus senegalensis	23.38	52	4.37 4557
Selene dorsalis	18.97	710	3.54
Galeoides decadactylus	15.70	151	2.93 4556
Rhizoprionodon acutus	14.48	4	2.70
Sardinella maderensis	8.05	120	1.50 4554
Pteroscion peli	6.66	217	1.24
Trichiurus lepturus	6.37	188	1.19
Chloroscombrus chrysurus	5.94	87	1.11
Penaeus notialis	5.28	250	0.99 4560
Portunus validus	4.49	23	0.84
Alectis alexandrinus	3.02	2	0.56
Caranx senegallus	1.97	2	0.37
Stromateus fiatola	1.90	4	0.35
Sardinella aurita	1.51	2	0.28
Caranx crysos	1.43	6	0.27
Scomberomorus tritor	1.37	2	0.26
Eucinostomus melanopterus	1.30	29	0.24
Pseudotolithus senegalensis	1.30	14	0.24
Balistes capricrus	0.62	2	0.12
Alectis alexandrinus	0.29	29	0.05
Total	535.35	99.98	

PROJECT STATION:1055
DATE:10/ 5/06 GEAR TYPE: PT No: 7 POSITION:Lat N 716
start stop duration Long W 1218
TIME :03:07:17 03:37:19 30 (min) Purpose code: 1
LOG :8049.75 8051.24 1.48 Area code : 3
FDEPTH: 12 14 GearCond.code:
BDEPTH: 24 32 Validity code:
Towing dir: 215° Wire out: 130 m Speed: 30 kn*10

Sorted: 73 Kg Total catch: 73.99 CATCH/HOUR: 147.98

SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
	weight	numbers	
Ilisha africana	63.80	3284	43.11 4537
Chloroscombrus chrysurus	25.50	676	17.23 4540
Sphyraena sphyraena	17.56	58	11.87 4538
Sardinella maderensis	14.64	214	9.89 4544
Galeoides decadactylus	6.70	90	4.53 4541
Sarda sarda	4.30		2.91
Alectis alexandrinus	4.20	48	2.84 4539
Brachydeuterus auritus	3.58	58	2.42 4543
Trichiurus lepturus	3.28		2.22
Drepane africana	1.20	2	0.81
Caranx crysos	0.60	4	0.41
Penaeus notialis	0.10	18	0.07
Pteroscion peli	0.10	2	0.07
Portunus validus	0.06	2	0.04
Penaeus kerathurus	0.04	2	0.03
Caranx hippos	0.04	2	0.03
Total	147.98	100.02	

PROJECT STATION:1059
DATE:10/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 710
start stop duration Long W 1222
TIME :09:40:22 10:10:28 30 (min) Purpose code: 3
LOG :8077.38 8078.99 1.60 Area code : 3
FDEPTH: 58 64 GearCond.code:
BDEPTH: 58 64 Validity code:
Towing dir: 215° Wire out: 180 m Speed: 30 kn*10

Sorted: 4 Kg Total catch: 4.91 CATCH/HOUR: 9.82

SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
	weight	numbers	
Scorpaena stephanica	3.58	8	36.46
Dentex angolensis	1.40	6	14.26
Priacanthus arenatus	1.16	8	11.81
Sphyraena guachancho	1.00	8	10.18
Aluterus heudeletii	0.56	2	5.70
Brachydeuterus auritus	0.50	8	5.09
Brotula barbata	0.46	2	4.68
Fistularia petimba	0.38	10	3.87
Pagrus caeruleostictus	0.38	2	3.87
Uranoscopus albusca	0.24	2	2.44
Selar crumenophthalmus	0.14	2	1.43
Pseudupeneus prayensis	0.02	2	0.20
Total	9.82	99.99	

PROJECT STATION:1060
DATE:10/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 648
start stop duration Long W 1148
TIME :15:32:31 16:02:51 30 (min) Purpose code: 3
LOG :0132.33 0133.90 1.55 Area code : 3
FDEPTH: 82 90 GearCond.code:
BDEPTH: 82 90 Validity code:
Towing dir: 219° Wire out: 225 m Speed: 30 kn*10

Sorted: 112 Kg Total catch: 4482.90 CATCH/HOUR: 8965.80

SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
	weight	numbers	
Ariommabondi	5962.80		66.51
Priacanthusarenatus	2627.40	84734	29.30 4562
Dentex congolensis	95.80	1020	1.07 4563
Sardinella maderensis	77.00	4086	0.86 4564
Sardinella aurita	70.00	2202	0.78 4565
Pagrus caeruleostictus	44.80	78	0.50
Decapterus rhonchus	29.80	630	0.33
Brachydeuterus auritus	27.52	1652	0.31
Fistularia petimba	12.58	78	0.14
Dentex angolensis	10.22		0.11
Decapterus punctatus	7.86	394	0.09
Total	8965.78	100.00	

PROJECT STATION:1061							
DATE:10/ 5/06	GEAR TYPE: BT No:19	POSITION:Lat N 654		Total	33.22	99.99	
start stop duration		Long W 1143					
TIME :16:41:00	16:41:00	30 (min)	Purpose code: 3				
LOG :8145.87	8147.34	1.46	Area code : 3				
FDEPTH: 45	50	GearCond.code:					
BDEPTH: 45	50	Validity code:					
Towing dir: 219°	Wire out: 150 m	Speed: 30 kn*10					
Sorted: 34 Kg	Total catch: 151.58	CATCH/HOUR: 303.16					
SPECIES	CATCH/HOUR	% OF TOT. C	SAMP				
	weight numbers						
Engraulis encrasicolus	140.80	7314	46.44				
Sphyraena guachancho	78.40	2584	25.86	4566			
Brachydeuterus auritus	45.28	8662	14.94				
Epinephelus aeneus	8.90	2	2.94	4570			
Sardinella maderensis	8.40	1280	2.77				
Pagrus caeruleostictus	3.80	10	1.25	4568			
Selene dorsalis	3.68	528	1.21				
Alectis alexandrinus	2.64	2	0.87				
Caranx cryos	2.42	4	0.80				
Selene dorsalis	2.32	14	0.77				
Sphyraena guachancho	1.98	18	0.65	4567			
Pseudotolithus senegalensis	1.74	2	0.57				
Galeoides decadactylus	1.44	72	0.47	4569			
Decapterus rhonchus	0.48	8	0.16				
Aluterus heudelotii	0.48	8	0.16				
Illex coindetii	0.40	104	0.13				
Total	303.16	99.99					
SPECIES	CATCH/HOUR	% OF TOT. C	SAMP				
	weight numbers						
Ilisha africana	168.00	4708	27.71	4603			
Pteroscion peli	82.80	3350	13.66				
Sphyraena guachancho	59.00	164	9.73	4597			
Pseudotolithus senegalensis	56.90	230	9.38	4598			
Trichiurus lepturus	42.40	700	6.99				
Parapenaeopsis atlantica	37.60	9520	6.20				
Pseudotolithus senegalensis	36.00	528	5.94	4600			
Etmalosa fimbriata	27.40	84	4.52	4599			
Chloroscombrus chrysurus	20.16	1200	3.32	4601			
Galeoides decadactylus	17.52	392	2.89	4602			
Brachydeuterus auritus	12.16	272	2.01				
Penaeus notialis	8.66	504	1.43	4606			
Pentanemus quinquarius	7.52	144	1.24	4604			
Cynoglossus senegalensis	5.44	48	0.90				
Pseudotolithus elongatus	5.20	64	0.86	4605			
Selene dorsalis	4.64		0.77				
Sardinella maderensis	3.20	312	0.53				
Sphyraena guachancho	2.72	32	0.45				
Echelus myrus	2.24	4	0.37				
Cynoglossus senegalensis	1.48	4	0.24				
Pseudotolithus elongatus	1.20	6	0.20				
Drepane africana	1.12	24	0.18				
Portunus validus	0.72	32	0.12				
Ephippion guttifer	0.64	16	0.11				
Lagocephalus laevigatus	0.64	8	0.11				
Decapterus rhonchus	0.40	8	0.07				
Pomadasys jubelini	0.32	8	0.05				
Brotula barbata	0.24	40	0.04				
Total	606.32	100.02					
SPECIES	CATCH/HOUR	% OF TOT. C	SAMP				
	weight numbers						
Parapenaeopsis atlantica	51.12	21.00					
Brachydeuterus auritus	44.32	1916	18.21	4577			
Pseudotolithus senegalensis	21.12	312	8.68	4572			
Pteroscion peli	20.00	822	8.22	4573			
Selene dorsalis	19.28	2348	7.92	4578			
Pseudotolithus senegalensis	19.10	110	7.85	4571			
Chloroscombrus chrysurus	15.20	1668	6.24	4580			
Pentanemus quinquarius	13.04	224	5.36	4582			
Ilisha africana	11.60	644	4.77	4574			
Trichiurus lepturus	5.92	96	2.43	4581			
Cynoglossus senegalensis	5.68	22	2.33	4576			
Penaeus notialis	2.60	114	1.07	4583			
Galeoides decadactylus	2.40	56	0.99	4579			
Portunus validus	2.22	6	0.91				
Sphyraena guachancho	2.14	6	0.88				
Drepane africana	2.12	2	0.87				
Callinectes pallidus	2.08	104	0.85				
Cynoglossus senegalensis	1.28	56	0.53	4575			
Sardinella maderensis	0.88	48	0.36				
B I V A L V E S	0.48	48	0.20				
Drepane africana	0.42	16	0.17				
Cynoglossus monodi	0.32	8	0.13				
Total	243.32	99.97					
Total	606.32	100.02					
SPECIES	CATCH/HOUR	% OF TOT. C	SAMP				
	weight numbers						
Parapenaeopsis atlantica	51.12	21.00					
Brachydeuterus auritus	44.32	1916	18.21	4577			
Pteroscion peli	20.00	822	8.22	4573			
Selene dorsalis	19.28	2348	7.92	4578			
Pseudotolithus senegalensis	19.10	110	7.85	4571			
Chloroscombrus chrysurus	15.20	1668	6.24	4580			
Pentanemus quinquarius	13.04	224	5.36	4582			
Ilisha africana	11.60	644	4.77	4574			
Trichiurus lepturus	5.92	96	2.43	4581			
Cynoglossus senegalensis	5.68	22	2.33	4576			
Penaeus notialis	2.60	114	1.07	4583			
Galeoides decadactylus	2.40	56	0.99	4579			
Portunus validus	2.22	6	0.91				
Sphyraena guachancho	2.14	6	0.88				
Drepane africana	2.12	2	0.87				
Callinectes pallidus	2.08	104	0.85				
Cynoglossus senegalensis	1.28	56	0.53	4575			
Sardinella maderensis	0.88	48	0.36				
B I V A L V E S	0.48	48	0.20				
Drepane africana	0.42	16	0.17				
Cynoglossus monodi	0.32	8	0.13				
Total	243.32	99.97					
Total	606.32	100.02					
SPECIES	CATCH/HOUR	% OF TOT. C	SAMP				
	weight numbers						
Brachydeuterus auritus	56.20	712	66.06	4608			
Epinephelus aeneus	17.60	6	20.69	4606			
Sphyraena guachancho	2.94	24	3.46	4607			
Brotula barbata	2.08	2	2.44				
Selene dorsalis	1.34	48	1.57	4610			
Trichiurus lepturus	0.82	14	0.96				
Pteroscion peli	0.74	12	0.87				
Pagellus bellottii	0.74	4	0.87				
Sardinella maderensis	0.64	80	0.75	4609			
Lagocephalus laevigatus	0.58	4	0.68				
Parapenaeus longirostris	0.48	32	0.56				
Dentex angolensis	0.44	2	0.52				
Saurida brasiliensis	0.28	44	0.33				
Illex coindetii	0.12	22	0.14				
Cepola sp.	0.08	2	0.09				
Total	85.08	99.99					
SPECIES	CATCH/HOUR	% OF TOT. C	SAMP				
	weight numbers						
Brachydeuterus auritus	56.20	712	66.06	4608			
Epinephelus aeneus	17.60	6	20.69	4606			
Sphyraena guachancho	2.94	24	3.46	4607			
Brotula barbata	2.08	2	2.44				
Selene dorsalis	1.34	48	1.57	4610			
Trichiurus lepturus	0.82	14	0.96				
Pteroscion peli	0.74	12	0.87				
Pagellus bellottii	0.74	4	0.87				
Sardinella maderensis	0.64	80	0.75	4609			
Lagocephalus laevigatus	0.58	4	0.68				
Parapenaeus longirostris	0.48	32	0.56				
Dentex angolensis	0.44	2	0.52				
Saurida brasiliensis	0.28	44	0.33				
Illex coindetii	0.12	22	0.14				
Cepola sp.	0.08	2	0.09				
Total	85.08	99.99					
SPECIES	CATCH/HOUR	% OF TOT. C	SAMP				
	weight numbers						
Chloroscombrus chrysurus	57.40	1024	51.37	4590			
Ilisha africana	14.58	460	13.05	4586			
Lutjanus dentatus	12.50	2	11.19				
Sphyraena guachancho	10.76	96	9.63	4584			
Sardinella maderensis	3.46	46	3.10	4588			
Selene dorsalis	3.12	40	2.79	4585			
Scomberomorus tritor	2.94	12	2.63				
Sardinella maderensis	2.18	310	1.95	4589			
Brachydeuterus auritus	2.10	76	1.88	4587			
Trichiurus lepturus	2.08	34	1.86				
Galeoides decadactylus	0.28	2	0.25				
Portunus validus	0.16	10	0.14				
Pentanemus quinquarius	0.12	2	0.11				
Penaeus notialis	0.06	6	0.05				
Total	111.74	100.00					
Total	111.74	100.01					
SPECIES	CATCH/HOUR	% OF TOT. C	SAMP				
	weight numbers						
Chloroscombrus chrysurus	DATE:11/ 5/06	GEAR TYPE: PT No: 7 POSITION:Lat N 641					
TIME :03:44:07	04:14:03	30 (min)	Purpose code: 1				
LOG :8189.53	8191.16	1.63	Area code : 4				
FDEPTH: 18	36	GearCond.code:					
BDEPTH: 75	64	Validity code:					
Towing dir: 35°	Wire out: 130 m	Speed: 30 kn*10					
Sorted: 4 Kg	Total catch: 16.61	CATCH/HOUR: 33.22					
SPECIES	CATCH/HOUR	% OF TOT. C	SAMP				
	weight numbers						
Pteroscion peli	25.30	822	76.16				
Brachydeuterus auritus	2.06	104	6.20	4592			
Selene dorsalis	1.50	28	4.52	4595			
Sphyraena guachancho	1.30	18	3.91	4596			
Chloroscombrus chrysurus	1.18	22	3.55	4594			
Sardinella maderensis	0.84	8	2.53				
Sepia officinalis hierredda	0.32	2	0.96				
Ilisha africana	0.24	14	0.72	4591			
Sardinella maderensis	0.20	28	0.60	4593			
Ariomma bondi	0.14	6	0.42				
Engraulis encrasiculus	0.08	10	0.24				
Trichiurus lepturus	0.06	2	0.18				
Total	112.19	100.01					
Total	112.19	100.01					
SPECIES	CATCH/HOUR	% OF TOT. C	SAMP				
	weight numbers						
Chloroscombrus chrysurus	DATE:11/ 5						

SPECIES	CATCH/HOUR	% OF TOT.	C	SAMP
	weight	numbers		
Dentex congoensis	46.40	530	60.81	4616
Illex coindetii	5.96	350	7.81	4619
Pentheroscion mbizi	51.14	36	6.74	4617
Dentex angolensis	4.90	38	6.42	4615
Ariomma bondi	2.92	94	3.83	4620
Squatina oculata	2.78	4	3.64	
Scorpaena sp.	2.28	4	2.99	
Fistularia petimba	1.48	8	1.94	
Priacanthus arenatus	1.32	12	1.73	
Antigonus capros	0.94	16	1.23	
Alloteuthis africana	0.72	172	0.94	
Lagocephalus laevisgatus	0.34	2	0.45	
Pagellus bellottii	0.30	2	0.39	
Epinephelus aeneus	0.24	2	0.31	4618
Trichurus lepturus	0.20	4	0.26	
Zeus faber	0.20	4	0.26	
Lepidotrigla carolae	0.18	4	0.24	
Total	76.30	99.99		

Sardinella maderensis	27.93	18.08	
Chloroscombrus chrysurus	21.41	13.86	4645
Sphyraena guachancho	20.28	13.13	
Brachydeuterus auritus	6.41	4.15	4646
Arius heudelotii	4.99	2	3.23
Scomberomorus tritor	3.04	4	1.97
Trichurus lepturus	2.83	41	1.83
Galeoides decadactylus	2.54	1.64	
Pomadasys jubelini	1.92	2	1.24
Selene dorsalis	1.84		1.19
Sphyraena guachancho	1.10	41	0.71
Chloroscombrus chrysurus	1.06	27	0.69
Sardinella maderensis	0.43	4	0.28
Parapeneopis atlantica	0.39	137	0.25
Drepane africana	0.39	2	0.25
Selene dorsalis	0.31	8	0.20
Sardinella maderensis	0.27	31	0.17
Brachydeuterus auritus	0.27	41	0.17
Penaeus notialis	0.02	4	0.01
Total	154.43	99.96	

Total 76.30 99.99

PROJECT STATION:1069

DATE:11/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 620
 start stop duration Long W 1055
 TIME :18:05:32 18:36:12 31 (min) Purpose code: 3
 LOG :8289.15 8290.80 1.64 Area code : 4
 FDEPTH: 43 51 GearCond.code:
 BDEPTH: 43 51 Validity code:
 Towing dir: 217° Wire out: 160 m Speed: 30 kn*10

Sorted: 43 Kg Total catch: 104.69 CATCH/HOUR: 202.63

SPECIES	CATCH/HOUR	% OF TOT.	C	SAMP
	weight	numbers		
Brachydeuterus auritus	57.48	5307	28.37	4630
Sardinella maderensis	56.32	6288	27.79	4624
Sphyraena guachancho	13.82	331	6.82	4627
Selene dorsalis	11.15	2388	5.50	
Sphyraena guachancho	10.59	37	5.23	4625
Trichurus lepturus	8.83	159	4.36	
Engraulis encrasicolus	6.50	2775	3.21	4623
Pseudotolithus senegalensis	6.45	21	3.18	4626
Pentheroscion mbizi	4.35	35	2.15	
Penaeus notialis	3.12	182	1.54	4622
Ilisha africana	3.08	70	1.52	4628
Galeoides decadactylus	2.55	64	1.26	4629
Raja miraletus	2.21	6	1.09	
Pteroscion peli	2.09	87	1.03	
Brotula barbata	1.76	4	0.87	
Balistes capriscus	1.63	6	0.80	
Penaeus notialis	1.57	151	0.77	4621
Pentheroscion mbizi	1.57	10	0.77	
Illex coindetii	1.45	64	0.72	
Chloroscombrus chrysurus	1.10	12	0.54	
Sepia sp.	0.99	29	0.49	
Parapeneopis atlantica	0.99	105	0.49	
Panulirus regius	0.91	4	0.45	
Pseudotolithus senegalensis	0.81	6	0.40	
Sphyraena sphyraena	0.41	2	0.20	
Saurida brasiliensis	0.35	81	0.17	
Cynoglossus senegalensis	0.23	6	0.11	
Syacium micrum	0.17	12	0.08	
Alectis alexandrinus	0.12	6	0.06	
Penaeus kerathurus	0.04	2	0.02	
Total	202.64	99.99		

PROJECT STATION:1070

DATE:11/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 623
 start stop duration Long W 1052
 TIME :19:43:19 20:13:17 30 (min) Purpose code: 3
 LOG :8298.96 8300.56 1.58 Area code : 4
 FDEPTH: 20 29 GearCond.code:
 BDEPTH: 20 29 Validity code:
 Towing dir: 217° Wire out: 140 m Speed: 30 kn*10

Sorted: 53 Kg Total catch: 132.61 CATCH/HOUR: 265.22

SPECIES	CATCH/HOUR	% OF TOT.	C	SAMP
	weight	numbers		
Chloroscombrus chrysurus	34.08	844	12.85	4638
Pseudotolithus senegalensis	33.20	162	12.52	4632
Pseudotolithus senegalensis	32.16	674	12.13	4633
Parapeneopis atlantica	30.96	13336	11.67	
Pteroscion peli	28.16	1490	10.62	4642
Trichurus lepturus	20.88	522	7.87	
Brachydeuterus auritus	17.04	1242	6.42	4641
Selene dorsalis	11.92	1192	4.49	4639
Sardinella maderensis	10.88	564	4.10	4643
Sphyraena guachancho	7.96	26	3.00	4634
Ilisha africana	7.28	656	2.74	
Galeoides decadactylus	4.72	216	1.78	4640
Cynoglossus senegalensis	4.20	14	1.58	4636
Pentanemus quinquearius	3.92	88	1.48	4635
Pseudotolithus elongatus	3.92	8	1.48	4637
Pomadasys jubelini	2.52	2	0.95	
Drepane africana	1.68	96	0.63	
Penaeus notialis	1.68	46	0.63	4631
Sphyraena guachancho	1.52	40	0.57	
Ephippion guttifer	1.12	32	0.42	
Trachinotus ovatus	1.04	16	0.39	
Cynoglossus sp.	1.04	80	0.39	
Cynoglossus senegalensis	0.96	16	0.36	
Caranx hippos	0.92	4	0.35	
Antennarius occidentalis	0.80	72	0.30	
Etmalosa fimbriata	0.36	2	0.14	
Portunus validus	0.30	2	0.11	
Total	265.22	99.97		

PROJECT STATION:1071

DATE:12/ 5/06 GEAR TYPE: PT No: 7 POSITION:Lat N 600
 start stop duration Long W 1016
 TIME :03:07:57 03:37:13 29 (min) Purpose code: 1
 LOG :8347.15 8348.63 1.72 Area code : 4
 FDEPTH: 16 22 GearCond.code:
 BDEPTH: 28 35 Validity code:
 Towing dir: 215° Wire out: 130 m Speed: 30 kn*10

Sorted: 72 Kg Total catch: 74.65 CATCH/HOUR: 154.45

SPECIES	CATCH/HOUR	% OF TOT.	C	SAMP
	weight	numbers		
Ilisha africana	57.00	3209	36.91	4644

SPECIES	CATCH/HOUR	% OF TOT.	C	SAMP
	weight	numbers		
Alectis alexandrinus	1401.71	309	81.33	4647
Caranx senegallus	233.00		13.52	
Selene dorsalis	78.57	597	4.56	4648
Sardinella maderensis	8.06	57	0.47	4649
Ilisha africana	2.09	46	0.12	4650
Total	1723.43		100.00	

SPECIES	CATCH/HOUR	% OF TOT.	C	SAMP
	weight	numbers		
Drepane africana	18.41	106	36.31	4651
Sphyraena guachancho	14.32		28.24	
Galeoides decadactylus	4.01		7.91	
Pseudotolithus senegalensis	3.90		7.69	
Ephippion guttifer	3.00	3	5.92	
Panulirus regius	2.84	5	5.60	
Scomberomorus tritor	1.23	3	2.43	
Penaeus kerathurus	0.82	38	1.62	
Cynoglossus senegalensis	0.60	3	1.18	
Pomadasys jubelini	0.49	5	0.97	
Chaetodipterus goreensis	0.30	3	0.59	
Penaeus notialis	0.22	19	0.43	
Eucinostomus melanopterus	0.16	3	0.32	
Sardinella maderensis	0.16	3	0.32	
Pseudupeneus prayensis	0.16	3	0.32	
Alectis alexandrinus	0.08	3	0.16	
Total	50.70		100.01	

SPECIES	CATCH/HOUR	% OF TOT.	C	SAMP
	weight	numbers		
Drepane africana	24.40	8	30.36	4654
Selene dorsalis	15.20	290	18.91	4652
Dentex angolensis	15.20	90	18.91	4655
Alloteuthis africana	9.66	4186	12.02	
Fistularia petimba	2.46	20	3.06	
Epinephelus aeneus	2.28	2	2.84	
Brachydeuterus auritus	1.76	254	2.19	4656
Caranx senegallus	1.74	2	2.17	
Sphyraena guachancho	1.66	6	2.07	
Pagrus caeruleostictus	1.30	6	1.62	4657
Chloroscombrus chrysurus	0.86	10	1.07	
Decapterus rhonchus	0.78	6	0.97	4653
Boops boops	0.64		0.80	
Lagocephalus laevisgatus	0.64	2	0.80	
Pomadasys jubelini	0.58	2	0.72	
Saurida brasiliensis	0.48	88	0.60	
Galeoides decadactylus	0.30	2	0.37	
Pseudupeneus prayensis	0.22	4	0.27	
Ilisha africana	0.14	4	0.17	
Sardinella maderensis	0.06	4	0.07	
Total	80.36		99.99	

SPECIES	CATCH/HOUR	% OF TOT.	C	SAMP
	weight	numbers		
Alectis alexandrinus	20.52	8	25.63	4658
Caranx hippos	15.20	290	18.91	4652
Antennarius occidentalis	15.20	90	18.91	4655
Etmalosa fimbriata	9.66	4186	12.02	
Portunus validus	2.46	20	3.06	
Sorted: 65 Kg Total catch: 213.30 CATCH/HOUR: 426.60				

SPECIES	CATCH/HOUR	% OF TOT.	C	SAMP
	weight	numbers		
Drepane africana	20.52	8	25.63	4658
Sphyraena guachancho	15.20	290	18.91	4652
Brachydeuterus auritus	15.20	90	18.91	4655
Alloteuth				

Trachurus trecae	105.00	1274	24.61	4662
Dentex congolensis	89.60	1164	21.00	4659
Dentex angolensis	85.40	630	20.02	4663
Boops boops	53.40	478	12.52	4660
Priacanthus arenatus	53.20	932	12.47	4658
Dentex canariensis	13.20	18	3.09	4661
Scorpaena sp.	12.00	22	2.81	
Fistularia petimba	6.40	22	1.50	
Brotula barbata	2.80	8	0.66	
Anthias anthias	1.96	106	0.46	
Pagellus bellottii	1.68	8	0.39	
Pseudupeneus prayensis	0.98	8	0.23	
Lagocephalus laevigatus	0.84	8	0.20	
Ariomma bondi	0.14	8	0.03	

Total 426.60 99.99

Priacanthus arenatus	0.94	9	0.47
Torpedo torpedo	0.83	14	0.41
Dactylopterus volitans	0.66	6	0.33
Sepia officinalis hierredda	0.54	9	0.27
Chaetodipterus goorensis	0.49	6	0.24
Brotula barbata	0.20	6	0.10
Syacium micrum	0.20	6	0.10
Decapterus rhonchus	0.17	14	0.08

Total 201.54 100.03

PROJECT STATION:1079

DATE:13/ 5/06 GEAR TYPE: PT No: 7 POSITION:Lat N 503
start stop duration Long W 915
TIME :04:12:19 04:42:09 30 (min) Purpose code: 1
LOG :8510.63 8512.31 1.65 Area code : 4
FDEPTH: 18 35 GearCond.code:
BDEPTH: 59 64 Validity code:
Towing dir: 214° Wire out: 130 m Speed: 32 kn*10

Sorted: 2 Kg Total catch: 2.62 CATCH/HOUR: 5.24

PROJECT STATION:1076

DATE:12/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 521
start stop duration Long W 953
TIME :17:23:08 17:46:50 24 (min) Purpose code: 3
LOG :8445.83 8446.97 1.13 Area code : 4
FDEPTH: 87 93 GearCond.code:
BDEPTH: 87 93 Validity code:
Towing dir: 220° Wire out: 230 m Speed: 30 kn*10

Sorted: 100 Kg Total catch: 100.20 CATCH/HOUR: 250.50

SPECIES	CATCH/HOUR		% OF TOT. C	SAMP
	weight	numbers		
Dentex congolensis	165.50	2040	66.07	4668
Dentex angolensis	44.50	393	17.76	4666
Dentex canariensis	9.88	15	3.94	4669
Pagellus bellottii	9.75	78	3.89	4665
Ariomma bondi	7.68	308	3.07	4667
Priacanthus arenatus	4.23	70	1.69	
Scorpaena scrofa	2.80	8	1.12	
Lagocephalus laevigatus	1.85	13	0.74	
Fistularia petimba	1.20	15	0.48	
Boops boops	0.95	10	0.38	
Pseudupeneus prayensis	0.63	5	0.25	
Illex coindetii	0.60	38	0.24	
Lepidotrigla cadmani	0.40	10	0.16	
Chelidonichthys gabonensis	0.38	5	0.15	
Lepidotrigla carolae	0.20	10	0.08	

Total 250.55 100.02

SPECIES	CATCH/HOUR		% OF TOT. C	SAMP
	weight	numbers		
Raja miraletus	1.72	6	32.82	
Arius heudeleti	0.78	2	14.89	
Illex coindetii	0.76	10	14.50	4688
Engraulis encrasicolus	0.56	152	10.69	4687
Chloroscombrus chrysurus	0.46	12	8.78	4683
Decapterus punctatus	0.38	30	7.25	4684
Sardinella aurita	0.16	6	3.05	4685
Torpedo torpedo	0.12	2	2.29	
Citharus linguatula	0.10	2	1.91	
Ilisha africana	0.10	2	1.91	
Brachydeuterus auritus	0.06	14	1.15	4686
Penaeus notialis	0.02	2	0.38	
Boops boops	0.02	2	0.38	

Total 5.24 100.00

PROJECT STATION:1080

DATE:13/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 503
start stop duration Long W 915
TIME :06:17:32 06:47:39 30 (min) Purpose code: 3
LOG :8517.35 8518.95 1.58 Area code : 4
FDEPTH: 54 63 GearCond.code:
BDEPTH: 54 63 Validity code:
Towing dir: 215° Wire out: 180 m Speed: 30 kn*10

Sorted: 44 Kg Total catch: 44.06 CATCH/HOUR: 88.12

DATE:12/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 526
start stop duration Long W 948
TIME :19:05:55 19:35:38 30 (min) Purpose code: 3
LOG :8456.03 8457.72 1.68 Area code : 4
FDEPTH: 66 61 GearCond.code:
BDEPTH: 66 61 Validity code:
Towing dir: 35° Wire out: 200 m Speed: 30 kn*10

Sorted: 36 Kg Total catch: 107.92 CATCH/HOUR: 215.84

SPECIES	CATCH/HOUR		% OF TOT. C	SAMP
	weight	numbers		
Dentex angolensis	93.10	680	43.13	4673
Brachydeuterus auritus	47.60	3328	22.05	4671
Dactylopterus volitans	10.64	50	4.93	
Brotula barbata	7.84	20	3.63	4675
Saurida brasiliensis	7.00	966	3.24	
Sphyraena guachancho	6.94	12	3.22	4672
Cynoscion senegalensis	5.88	42	2.72	4677
Citharus linguatula	5.04	154	2.34	
Pseudupeneus prayensis	5.04	132	2.34	4670
Scorpaena sp.	4.74	8	2.20	
Ariomma bondi	4.00	98	1.85	
Pagellus bellottii	3.86	42	1.79	
Parapenaeus longirostris	2.52	608	1.17	
Umbrina canariensis	2.38	8	1.10	4676
Priacanthus arenatus	1.82	22	0.84	
Boops boops	1.54	246	0.71	
Illex coindetii	1.26	22	0.58	
Sepla officinalis hierredda	0.98	22	0.45	
Decapterus punctatus	0.98	98	0.45	4674
Uranoscopus albusca	0.92	8	0.43	
Physiculus huloti	0.50	36	0.23	
Arnoglossus imperialis	0.36	42	0.17	
Sardinella maderensis	0.28	36	0.13	
Lepidotrigla carolae	0.28	22	0.13	
Bathygobius paganelus	0.22	36	0.10	
MURAENESOCIDAE	0.22	14	0.10	

Total 215.94 100.03

SPECIES	CATCH/HOUR		% OF TOT. C	SAMP
	weight	numbers		
Selene dorsalis	25.50	404	28.94	4693
Pagrus caeruleostictus	17.70	42	20.09	4692
Epinephelus aeneus	13.90	2	15.77	
Pagellus bellottii	11.24	64	12.76	4694
Illex coindetii	4.76	56	5.40	
Dentex angolensis	4.44	28	5.04	4691
Torpedo torpedo	3.14	8	3.56	
Albulaa vulpes	1.48	2	1.68	
Sphyraena guachancho	1.18	30	1.34	
Pseudupeneus prayensis	0.98	32	1.11	4690
Umbrina canariensis	0.82	2	0.93	
Brachydeuterus auritus	0.80	60	0.91	
Sardinella aurita	0.56	38	0.64	4695
Decapterus punctatus	0.34	26	0.39	4689
Fistularia petimba	0.28	2	0.32	
Decapterus punctatus	0.28	2	0.32	
Ariomma bondi	0.26	4	0.30	
Boops boops	0.14	12	0.16	
Priacanthus arenatus	0.12	2	0.14	
Penaeus notialis	0.08	4	0.09	
Syacium micrum	0.08	2	0.09	
Saurida brasiliensis	0.04	10	0.05	

Total 88.12 100.03

PROJECT STATION:1081

DATE:13/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 503
start stop duration Long W 919
TIME :08:00:39 08:30:54 30 (min) Purpose code: 3
LOG :8525.75 8527.40 1.65 Area code : 4
FDEPTH: 71 73 GearCond.code:
BDEPTH: 71 73 Validity code:
Towing dir: 215° Wire out: 200 m Speed: 30 kn*10

Sorted: 85 Kg Total catch: 85.14 CATCH/HOUR: 170.28

DATE:12/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 530
start stop duration Long W 945
TIME :20:37:30 20:58:08 21 (min) Purpose code: 3
LOG :8463.26 8464.36 1.09 Area code : 4
FDEPTH: 40 47 GearCond.code:
BDEPTH: 40 47 Validity code:
Towing dir: 220° Wire out: 150 m Speed: 30 kn*10

Sorted: 47 Kg Total catch: 70.53 CATCH/HOUR: 201.51

SPECIES	CATCH/HOUR		% OF TOT. C	SAMP
	weight	numbers		
Pomadasys jubelini	44.57	57	22.12	4681
Penaeus notialis	29.83	1309	14.80	4683
Pteroscion peli	26.97	1080	13.38	4679
Pseudotolithus senegalensis	23.17	129	11.50	4678
Galeoides decadactylus	14.83	377	7.36	4682
Brachydeuterus auritus	12.34	2869	6.12	4680
Raja miraletus	7.94	17	3.94	
Monochirius hispidus	4.69	31	2.33	
Sphyraena guachancho	4.69	11	2.33	
Stromateus fairola	4.29	9	2.13	
Portunus validus	3.31	6	1.64	
Chromis sp.	2.66	360	1.32	
Ilisha africana	2.31	94	1.15	
Dasyatis margarita	2.03	6	1.01	
Pomadasys incisus	1.69	6	0.84	
Cynoscion senegalensis	1.69	17	0.84	
Paronchelis stauchi	1.63	411	0.81	
Trichiurus lepturus	1.46	14	0.72	
Saurida brasiliensis	1.43	177	0.71	
Illex coindetii	1.29	17	0.64	
Selene dorsalis	1.26	69	0.63	
Pomadasys peroteti	1.20	6	0.60	
Umbrina canariensis	1.20	3	0.60	
Grammoplites gruveli	1.03	69	0.51	

Total 170.28 99.98

PROJECT STATION:1082

DATE:13/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 450
start stop duration Long W 924
TIME :09:38:52 10:08:54 30 (min) Purpose code: 3
LOG :8535.25 8536.83 1.57 Area code : 4
FDEPTH: 92 85 GearCond.code:
BDEPTH: 92 85 Validity code:
Towing dir: 35° Wire out: 250 m Speed: 30 kn*10

Sorted: 66 Kg Total catch: 641.07 CATCH/HOUR: 1282.14

SPECIES CATCH/HOUR % OF TOT. C SAMP

	weight	numbers		
Trachurus trecae	301.50	6300	23.52	4710
Dentex congensis	284.40	3462	22.18	4704
Boops boops	202.50	2818	15.79	4708
Sardinella aurita	178.20	3794	13.90	4709
Dentex angolensis	51.30	486	4.00	4705
Decapterus punctatus	50.94	1152	3.97	4707
Pagellus bellottii	49.50	468	3.86	4711
Squatina oculata	31.68	18	2.47	
Priacanthus arenatus	28.08	846	2.19	4712
Scorpaena scrofa	23.40	54	1.83	
Umbrina canariensis	23.40	126	1.83	4706
Raja miraletus	8.64	18	0.67	
Pseudupeneus prayensis	7.56	72	0.59	
Scomber japonicus	7.20	126	0.56	
Dentex canariensis	7.02	18	0.55	
Sargocentron hastatum	5.22	36	0.41	
Ariomma bondi	4.86	108	0.38	
Lagocephalus laevisgatus	4.32	54	0.34	
Anthias anthias	4.14	342	0.32	
Syacium micrum	1.62	18	0.13	
Lepidotrigla cadmani	1.44	36	0.11	
Illex coindetii	1.44	18	0.11	
Fistularia petimba	1.44	36	0.11	
Sepia officinalis hierredda	1.26	18	0.10	
Chaetodon marcellae	1.08	18	0.08	

SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
Priacanthus arenatus	525.60	25562	48.13
Trachurus trecae	227.70	9056	20.85
Ariomma bondi	105.30	3346	9.64
Boops boops	91.44	4932	8.37
Dentex angolensis	32.40	54	2.97
Sphyraena guachancho	19.00	92	1.74
Pagellus bellottii	13.86	450	1.27
Sphyraena guachancho	12.78	72	1.17
Dentex congensis	10.36	42	0.95
Decapterus punctatus	8.64	270	0.79
Dentex angolensis	6.94	112	0.64
Scomber japonicus	6.28	6	0.58
Scorpaena scrofa	6.12	126	0.56
Lagocephalus laevisgatus	5.30	12	0.49
Raja miraletus	4.50	18	0.41
Pentheroscion mbizi	4.44	16	0.41
Pagellus bellottii	3.00	14	0.27
Sardinella aurita	2.40	50	0.22
Illex coindetii	2.16	54	0.20
Fistularia petimba	1.80	18	0.16
Zeus faber	0.78	4	0.07
Pseudupeneus prayensis	0.68	4	0.06
Total	1092.08	100.00	

Total 1282.14 100.00

PROJECT STATION:1083
DATE:13/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 432
start stop duration Long W 845
TIME :17:25:30 17:55:11 30 (min) Purpose code: 3
LOG :8606.48 8608.03 1.54 Area code : 4
FDEPTH: 85 99 GearCond.code:
BDEPTH: 85 99 Validity code:
Towing dir: 214° Wire out: 250 m Speed: 30 kn*10

Sorted: 104 Kg Total catch: 104.79 CATCH/HOUR: 209.58

PROJECT STATION:1086
DATE:14/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 413
start stop duration Long W 733
TIME :11:57:09 12:27:09 30 (min) Purpose code: 3
LOG :8732.98 8734.49 1.51 Area code : 4
FDEPTH: 77 80 GearCond.code:
BDEPTH: 77 80 Validity code:
Towing dir: 260° Wire out: 200 m Speed: 30 kn*10

Sorted: 73 Kg Total catch: 73.10 CATCH/HOUR: 146.20

SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
	weight	numbers	
Squatina oculata	90.00	2	42.94
Trachurus trecae	63.00	1014	30.06
Dentex congensis	39.30	502	18.75
Dentex angolensis	25.50	180	12.17
Boops boops	24.12	368	11.51
Dentex canariensis	20.00	14	9.54
Pagellus bellottii	7.70	90	3.67
Torpedo torpedo	5.00	2	2.39
Scomber japonicus	3.48	36	1.66
Scorpaena scrofa	2.60	4	1.24
Scorpaena angolensis	2.46	4	1.17
Zeus faber	1.50	6	0.72
Priacanthus arenatus	1.24	16	0.59
Illex coindetii	1.22	16	0.58
Umbrina canariensis	1.16	4	0.55
Fistularia petimba	0.64	4	0.31
Raja miraletus	0.56	2	0.27
Chilomycterus spinosus mauret.	0.40	2	0.19
Ariomma bondi	0.24	4	0.11
Lagocephalus laevisgatus	0.22	2	0.10
Pseudupeneus prayensis	0.14	2	0.07
Lepidotrigla cadmani	0.10	2	0.05

SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
	weight	numbers	
Sphyraena guachancho	77.20	1384	52.80
Selene dorsalis	11.84	320	8.10
Pentheroscion mbizi	11.70	82	8.00
Priacanthus arenatus	11.14	302	7.62
Sphyraena guachancho	10.78	58	7.37
Decapterus punctatus	4.64	130	3.17
Trichiurus lepturus	3.50	64	2.39
Sepia officinalis hierredda	3.38	10	2.31
Dentex angolensis	2.96	28	2.02
Illex coindetii	2.40	154	1.64
Sardinella aurita	1.44	50	0.98
Engraulis encrasicolus	1.26	138	0.86
Chelidonichthys gabonensis	0.86	10	0.59
Caranx cryos	0.84	4	0.57
Illex coindetii	0.46	6	0.31
Ariomma bondi	0.38	4	0.26
Citharus linguatula	0.28	4	0.19
Pseudupeneus prayensis	0.22	2	0.15
Dentex congensis	0.22	4	0.15
Lagocephalus laevisgatus	0.18	4	0.12
Fistularia petimba	0.16	6	0.11
Parapeneus longirostris	0.14	22	0.10
Bathygobius paganeus	0.04	2	0.03

Total 290.58 138.64

Total 146.02 99.84

PROJECT STATION:1084
DATE:13/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 437
start stop duration Long W 841
TIME :18:56:46 19:26:45 30 (min) Purpose code: 3
LOG :8615.92 8617.55 1.62 Area code : 4
FDEPTH: 73 69 GearCond.code:
BDEPTH: 73 69 Validity code:
Towing dir: 35° Wire out: 200 m Speed: 30 kn*10

Sorted: 80 Kg Total catch: 80.63 CATCH/HOUR: 161.26

PROJECT STATION:1087
DATE:14/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 417
start stop duration Long W 733
TIME :13:39:07 14:09:07 30 (min) Purpose code: 3
LOG :8742.30 8743.86 1.53 Area code : 4
FDEPTH: 38 40 GearCond.code:
BDEPTH: 38 40 Validity code:
Towing dir: 265° Wire out: 150 m Speed: 30 kn*10

Sorted: 112 Kg Total catch: 277.68 CATCH/HOUR: 555.36

SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
	weight	numbers	
Dentex angolensis	54.00	630	33.49
Pagellus bellottii	22.60	284	14.01
Epinephelus aeneus	15.80	2	9.80
Ariomma bondi	10.36	220	6.42
Brachydeuterus auritus	7.18	126	4.45
Priacanthus arenatus	6.78	206	4.20
Chelidonichthys gabonensis	6.12	210	3.80
Chelidonichthys gabonensis	4.98	88	3.09
Illex coindetii	4.40	52	2.73
Brotula barbata	3.36	6	2.08
Branchiostegus semifasciatus	3.16	16	1.96
Citharus linguatula	3.06	182	1.90
Trachurus trecae	2.46	52	1.53
Umbrina canariensis	2.16	10	1.34
Cynoglossus senegalensis	1.72	10	1.07
Pseudupeneus prayensis	1.40	32	0.87
Mustelus mustelus	1.36	2	0.84
Sepia officinalis hierredda	1.30	26	0.81
Raja miraletus	1.28	6	0.79
Boops boops	1.22	28	0.76
NETTASTOMATIDAE	1.06	88	0.66
Sphyraena guachancho	0.76	2	0.47
Chilomycterus spinosus mauret.	0.70	8	0.43
Dactylopterus volitans	0.60	10	0.37
Argoneness imperialis	0.52	56	0.32
Pagrus caeruleostictus	0.46	2	0.29
Octopus vulgaris	0.36	2	0.22
Fistularia petimba	0.26	6	0.16
Uranoscopus cadenati	0.20	2	0.12
Scorpaena angolensis	0.20	2	0.12
Portunus validus	0.16	6	0.10
Sardinella aurita	0.10	10	0.06
Sphyraena guachancho	0.10	2	0.06

SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
	weight	numbers	
Sphyraena guachancho	210.10	642	37.83
Sphyraena guachancho	121.00	910	21.79
Galeoides decadactylus	52.20	158	9.40
Pseudotolithus senegalensis	47.10	140	8.48
Selene dorsalis	44.60	200	8.03
Brachydeuterus auritus	16.56	506	2.98
Stromateus fiatola	15.70	10	2.83
Chloroscombrus chrysurus	13.40	184	2.41
Sphyraena guachancho	10.68	60	1.92
Raja miraletus	3.70	6	0.67
Pseudotolithus senegalensis	3.36	40	0.61
Ilisha africana	2.90	82	0.52
Mustelus mustelus	2.26	2	0.41
Sardinella maderensis	1.76	130	0.32
Galeoides decadactylus	1.66	16	0.30
Pentheroscion mbizi	1.48	8	0.27
Cynoglossus canariensis	1.46	2	0.26
Portunus validus	1.32	8	0.24
Pomadasys jubelini	0.98	6	0.18
Selene dorsalis	0.78	76	0.14
Trichiurus lepturus	0.70	18	0.13
Elagatis bipinnulata	0.62	12	0.11
Scomberomorus tritor	0.58	2	0.10
Parapeneopeis atlantica	0.28	36	0.05
Fistularia petimba	0.16	2	0.03
Brotula barbata	0.02	2	

Total 555.36 100.01

PROJECT STATION:1085
DATE:14/ 5/06 GEAR TYPE: BT No:19 POSITION:Lat N 421
start stop duration Long W 803
TIME :06:20:29 06:50:14 30 (min) Purpose code: 3
LOG :8686.45 8687.96 1.49 Area code : 4
FDEPTH: 78 82 GearCond.code:
BDEPTH: 78 82 Validity code:
Towing dir: 215° Wire out: 200 m Speed: 30 kn*10


Sorted: 73 Kg Total catch: 546.04 CATCH/HOUR: 1092.08

SPECIES	CATCH/HOUR	% OF TOT. C	SAMP
	weight	numbers	
Pseudotolithus typus	634.66	131	58.66
Pseudotolithus typus	256.55	632	23.71
Arius heudeleti	48.00	14	4.44
Lutjanus dentatus	46.06	4	4.26
Selene dorsalis	28.59	131	2.64


Total 306.52 1081.84

<i>Galeoides decadactylus</i>	24.64	78	2.28	4759
<i>Umbrina canariensis</i>	14.65	7	1.35	4762
<i>Pseudotolithus senegalensis</i>	8.68	4	0.80	
<i>Chloroscombrus chrysurus</i>	4.55	32	0.42	
<i>Dasyatis margarita</i>	3.49	4	0.32	
<i>Pseudotolithus elongatus</i>	3.39	4	0.31	
<i>Stromateus fiatola</i>	2.93	7	0.27	
<i>Raja miraletus</i>	1.84	4	0.17	
<i>Pentanemus quinquarius</i>	1.76	14	0.16	
<i>Pomadasys rogeri</i>	1.41	4	0.13	
<i>Parapenaeopsis atlantica</i>	0.64	95	0.06	
Total	1081.84		99.98	


Annex II Length distributions of main species




Dentex angolensis GUINEA BISSAU
Mean length = 21.1 cm N = 69




Pseudupeneus prayensis GUINEA BISSAU
Mean length = 20.0 cm N = 174




Dentex congoensis GUINEA BISSAU
Mean length = 19.1 cm N = 125




Spicara alta GUINEA BISSAU
Mean length = 23.7 cm N = 85




Pagellus bellottii GUINEA BISSAU
Mean length = 21.7 cm N = 152




Brachydeuterus auritus GUINEA BISSAU
Mean length = 15.1 cm N = 128




Pagrus caeruleostictus GUINEA BISSAU
Mean length = 25.8 cm N = 39




Sardinella aurita GUINEA BISSAU
Mean length = 12.7 cm N = 194




Chloroscombrus chrysurus GUINEA BISSAU
Mean length = 18.7 cm N = 168




Scomber japonicus GUINEA BISSAU
Mean length = 16.5 cm N = 198




Decapterus punctatus GUINEA BISSAU
Mean length = 12.9 cm N = 202




Sepia officinalis hierredda GUINEA BISSAU
Mean length = 12.0 cm N = 49




Decapterus rhonchus GUINEA BISSAU
Mean length = 20.8 cm N = 325




Chlorophthalmus atlanticus GUINEA BISSAU
Mean length = 16.5 cm N = 93




Trachurus trecae GUINEA BISSAU
Mean length = 16.1 cm N = 247




Ariomma bondi GUINEA BISSAU
Mean length = 13.9 cm N = 89




Pagellus bellottii
GUINEA
Mean length = 20.5 cm
N = 111




Pseudupeneus prayensis
GUINEA
Mean length = 13.0 cm
N = 338




Pagrus caeruleostictus
GUINEA
Mean length = 12.1 cm
N = 355




Merluccius polli
GUINEA
Mean length = 32.9 cm
N = 97




Brachydeuterus auritus
GUINEA
Mean length = 13.1 cm
N = 111




Sardinella aurita
GUINEA
Mean length = 12.6 cm
N = 230



Lethrinus atlanticus
GUINEA
Mean length = 34.3 cm
N = 63




Sardinella maderensis
GUINEA
Mean length = 9.8 cm
N = 204




Engraulis encrasiculus
Mean length = 9.1 cm

GUINEA
N = 36




Trachurus trecae
Mean length = 12.0 cm

GUINEA
N = 179




Chloroscombrus chrysurus
Mean length = 18.0 cm

GUINEA
N = 135




Sphyraena guachancho
Mean length = 35.9 cm

GUINEA
N = 54




Decapterus punctatus
Mean length = 13.7 cm

GUINEA
N = 222




Dentex angolensis
Mean length = 20.7 mm

SIERRA LEONE
N = 158




Decapterus rhonchus
Mean length = 11.3 cm

GUINEA
N = 200




Dentex congoensis
Mean length = 17.7 cm


SIERRA LEONE
N = 268




Pagellus bellottii SIERRA LEONE
Mean length = 18.1 cm N = 715




Pseudotolithus elongatus SIERRA LEONE
Mean length = 33.2 cm N = 33




Pagrus caeruleostictus SIERRA LEONE
Mean length = 20.3 cm N = 885




Pseudotolithus senegalensis SIERRA LEONE
Mean length = 23.3 cm N = 222




Brachydeuterus auritus SIERRA LEONE
Mean length = 12.0 cm N = 521




Tethrinus atlanticus SIERRA LEONE
Mean length = 27.9 cm N = 68




Pomadasys rogeri SIERRA LEONE
Mean length = 41.6 cm N = 81




Pseudupeneus prayensis SIERRA LEONE
Mean length = 14.7 cm N = 452




Galeoides decadactylus
SIERRA LEONE
Mean length = 14.4 cm
N = 547




Sardinella aurita
SIERRA LEONE
Mean length = 11.8 cm
N = 386




Pantanemus quinquatus
SIERRA LEONE
Mean length = 18.2 cm
N = 122




Sardinella maderensis
SIERRA LEONE
Mean length = 12.8 cm
N = 580




Balistes capriscus
SIERRA LEONE
Mean length = 37.7 cm
N = 96




Ilisha africana
SIERRA LEONE
Mean length = 12.4 cm
N = 162




Psettodes sp.
SIERRA LEONE
Mean length = 28.0 cm
N = 74




Chloscombeus chrysurus
SIERRA LEONE
Mean length = 17.2 cm
N = 896




Decapterus punctatus
SIERRA LEONE
Mean length = 12.9 cm
N = 463




Sphyraena guachancho
SIERRA LEONE
Mean length = 21.2 cm
N = 707




Decapterus rhonchus
SIERRA LEONE
Mean length = 19.9 cm
N = 336




Ariommabondi
SIERRA LEONE
Mean length = 15.9 cm
N = 128




Selene dorsalis
SIERRA LEONE
Mean length = 10.2 cm
N = 147




Priacanthus arenatus
SIERRA LEONE
Mean length = 12.5 cm
N = 223




Trachurus trecae
SIERRA LEONE
Mean length = 17.3 cm
N = 108




Sepia officinalis hierredda
SIERRA LEONE
Mean length = 17.5 cm
N = 11




Penaeus notialis
Mean length = 32.4 mm
SIERRA LEONE
N = 356




Dentex canariensis
Mean length = 37.0 cm
LIBERIA
N = 37




Penaeus kerathurus
Mean length = 32.2 mm
SIERRA LEONE
N = 97




Dentex congoensis
Mean length = 17.5 cm
LIBERIA
N = 674




Boops boops
Mean length = 15.0 cm
LIBERIA
N = 201




Pagellus bellottii
Mean length = 18.7 cm
LIBERIA
N = 362




Dentex angolensis
Mean length = 19.4 cm
LIBERIA
N = 1123




Pagrus caeruleostictus
Mean length = 28.0 cm
LIBERIA
N = 29




Pseudotolithus senegalensis
LIBERIA
Mean length = 21.9 cm
N = 369




Umbrina canariensis
LIBERIA
Mean length = 25.2 cm
N = 17




Pseudotolithus typus
LIBERIA
Mean length 46.9 cm
N = 73




Brachydeuterus auritus
LIBERIA
Mean length = 9.2 cm
N = 704




Pentheroscion mbizi
LIBERIA
Mean length = 22.8 cm
N = 107




Pseudupeneus prayensis
LIBERIA
Mean length = 14.1 cm
N = 76




Pteroscion peli
LIBERIA
Mean length = 11.5 cm
N = 72




Galeoides decadactylus
LIBERIA
Mean length = 16.9 cm
N = 242




Pentanemus quinquearius
Mean length = 19.1 cm LIBERIA N = 29




Ethmalosa fimbriata
Mean length = 31.6 cm LIBERIA N = 42




Priacanthus arenatus
Mean length = 11.4 cm LIBERIA N = 398




Ilisha africana
Mean length = 14.4 cm LIBERIA N = 345




Priacanthus arenatus
Mean length = 17.7 cm LIBERIA N = 39




Sardinella aurita
Mean length = 17.0 cm LIBERIA N = 193




Ariommabondi
Mean length = 13.2 cm LIBERIA N = 367




Sardinella maderensis
Mean length = 10.2 cm LIBERIA N = 411




Engraulis encrasiculus
Mean length = 6.8 cm
LIBERIA
N = 102




Selene dorsalis
Mean length = 15.3 cm
LIBERIA
N = 367




Alectis alexandrinus
Mean length = 62.1 cm
LIBERIA
N = 112




Trachurus trecae
Mean length = 15.2 cm
LIBERIA
N = 346




Chloroscombrus chrysurus
Mean length = 14.8cm
LIBERIA
N = 247



Sphyraena sphyraena
Mean length = 27.1 cm
LIBERIA
N = 711



Decapterus punctatus
Mean length = 14.9 cm
LIBERIA
N = 288



Penaeus notialis
Mean length = 29.8 mm
LIBERIA
N = 285

Annex III Summary of biological samples

Species	Guinea-Bissau			Guinea			Sierra Leone			Liberia			Total		
	Min len	Max len	n meas	Min len	Max len	n meas	Min len	Max len	n meas	Min len	Max len	n meas	Min len	Max len	n meas
<i>Ethmalosa fimbriata</i>										28.1	35.0	20	28.1	35.0	20
<i>Sardinella aurita</i>	17.0	19.5	20	14.5	23.0	20							14.5	23.0	40
<i>Sardinella maderensis</i>				15.5	21.0	20				13.0	25.0	20	13.0	25.0	40
<i>Decapterus rhonchus</i>	15.0	33.0	27	17.0	31.5	18	22.0	26.0	20				15.0	33.0	65
<i>Trachurus trecae</i>	19.0	26.5	5				11.0	18.5	19				11.0	26.5	24
<i>Scomber japonicus</i>	12.5	20.5	20										12.5	20.5	20
<i>Sphyraena gauchancho</i>				24.0	55.0	20	22.0	52.0	20	18.5	44.0	20	18.5	55.0	60
<i>Priacanthus arenatus</i>	18.0	27.0	25										18.0	27.0	25
<i>Brachydeuterus auritus</i>	14.5	20.5	6										14.5	20.5	6
<i>Pomadasys jubelini</i>										32.0	50.0	13	32.0	50.0	13
<i>Pomadasys rogeri</i>							26.0	49.0	20				26.0	49.0	20
<i>Dentex angolensis</i>	16.0	21.5	24				18.0	26.5	20	15.5	23.0	20	15.5	26.5	64
<i>Dentex congogensis</i>	14.4	22.5	45				10.5	17.0	13	16.5	22.0	20	10.5	22.5	78
<i>Dentex macrophthalmus</i>	20.0	26.5	14										20.0	26.5	14
<i>Pagellus bellottii</i>				16.0	22.5	20	16.5	24.5	40				16.0	24.5	60
<i>Pagrus africanus</i>	20.5	43.0	6										20.5	43.0	6
<i>Pagrus caeruleostictus</i>	15.5	27.5	20	15.0	28.0	20	17.5	24.0	20				15.0	28.0	60
<i>Spicara alta</i>	17.5	25.0	20										17.5	25.0	20
<i>Lethrinus atlanticus</i>				27.5	40.0	20							27.5	40.0	20
<i>Pentheroscion mbizi</i>										18.5	30.5	20	18.5	30.5	20
<i>Pseudotolithus elongatus</i>							26.0	33.0	10				26.0	33.0	10
<i>Pseudotolithus senegalensis</i>							24.5	52.0	20	24.0	36.0	20	24.0	52.0	40
<i>Pseudupeneus prayensis</i>	18.5	27.0	20	11.0	24.5	20							11.0	27.0	40
<i>Galeoides decadactylus</i>							14.5	37.5	20				14.5	37.5	20

Annex IV Families/genera in catch analysis and swept area estimates

1) Main groups in swept-area bottom trawl hauls:

Demersal:

Sciaenidae, Sparidae, Haemulidae, Ariidae, Serranidae, Lutjanidae,
Merluccidae, Ophididae, Lethrinidae

Pelagic:

Carangidae, Scombridae, Sphyraenidae, Trichiuridae, Clupeidae, Engraulididae

Shrimp:

Shrimps

Cephalopods:

Cephalopods

Sharks:

Sharks

2) Main pelagic families in swept-area bottom trawl hauls:

Clupeids:

Clupeidae, Engraulididae

Carangids:

Carangidae

Scombrids:

Scombridae

Hairtails:

Trichiuridae

Barracudas:

Sphyraenidae

3) Commercially important demersal species grouped by families in swept-area bottom trawl hauls:

Seabream:

Dentex spp., *Diplodus* spp., *Lithognathus* spp., *Pagellus* spp., *Pagrus* spp., *Sparus* spp.

Snappers:

Lutjanidae

Groupers:

Serranidae

Grunts:

Plectorrhynchus spp., *Pomadasys* spp.

Croakers:

Sciaenidae

Annex V Swept area biomass estimates

SWEPT AREA ANALYSIS FROM STATION 987 TO STATION 1063

Guinea Bissau 2006

ONLY STATIONS IN SECTOR
ARE INCLUDED

SPECIES NAME	SAMPLE	DISTRIB. BY CATCH CLASSES					% inci-dence	Mean dens.t/nm ²	Mean densities by bottom depth strata t/nm ²			
		Lower limits, Kg/nm							- 30m	30- 50m	50-100m	100-200m
		>0	10	30	100	300	1000					
<i>Chlorophthalmus atlanticus</i>	1		1	1				18	3.90			
<i>Chloroscombrus chrysurus</i>	2		1	1	1			29	3.27	11.78	5.05	
<i>Decapterus rhonchus</i>	4	1	5	1				65	2.56	5.32	4.57	0.20
<i>Trachurus trecae</i>	5		2	1				47	2.20	0.02	6.08	1.33
<i>Pentheroscion mbizi</i>	2				1			18	2.02			8.57
<i>Pagellus bellottii</i>	5		1	1				41	1.83	8.76	1.13	0.06
<i>Brachydeuterus auritus</i>	1	1		2				18	1.73	0.79	6.77	
<i>Scomber japonicus</i>	3		3					35	1.30	3.30	2.20	0.56
<i>Spicara alta</i>	1		1	1				18	1.28			5.42
<i>Sardinella aurita</i>	4	1		1				35	0.97	4.36	0.64	0.13
<i>Todaropsis eblanae</i>	4	1	1					35	0.59			2.12
<i>Pseudupeneus prayensis</i>	5	2	1					47	0.45	1.14	0.92	0.08
<i>Antigonion capros</i>	4		1					29	0.39			0.05
<i>Pagrus caeruleostictus</i>	2		1					18	0.33	1.85	0.01	
<i>Dentex congensis</i>	2		1					12	0.33			1.14
<i>Scorpaena stephanica</i>	4	2						35	0.27			0.80
<i>Decapterus punctatus</i>	5	2						41	0.25	0.40	0.72	0.03
<i>Arius parkii</i>	2		1					18	0.24		1.03	
<i>Dasyatis pastinaca</i>				1				6	0.23		0.96	
<i>Synagrops microlepis</i>	3		1					12	0.23			0.98
<i>Dentex angolensis</i>	3	1						24	0.22			0.01
<i>Ariomma bondi</i>	4	1						29	0.20			0.36
<i>Rhizoprionodon acutus</i>	2	1						18	0.16	0.21	0.52	0.29
<i>Fistularia petimba</i>	9							41	0.14		0.33	0.11
<i>Mustelus mustelus</i>	2							12	0.11		0.15	0.24
<i>Sphoeroides marmoratus</i>	2	1						18	0.07		0.01	0.29
<i>Pagrus africanus</i>	1	1						12	0.07		0.01	0.27
<i>Caranx cryos</i>	4							24	0.07	0.32		0.03
<i>Albula vulpes</i>			1					6	0.07	0.40		
<i>Zenopsis conchifer</i>	2							12	0.06			0.25
<i>Sepia officinalis hierredda</i>	9							53	0.06	0.07	0.08	0.07
<i>Sphyraena afra</i>	1							6	0.06	0.31		
<i>Pseudotolithus senegalensis</i>	1							6	0.06		0.24	
<i>Zeus faber</i>	4							24	0.05		0.03	0.16
<i>Lagocephalus laevigatus</i>	7							41	0.05	0.02	0.08	0.07
<i>Saurida brasiliensis</i>	2							12	0.05			0.20
<i>Umbrina canariensis</i>	3							18	0.05			0.23
<i>Raja miraletus</i>	8							47	0.05		0.03	0.05
<i>Penaeus notialis</i>	1							6			0.96	0.46
Other fish									0.74	0.70		0.97
Sum all species								26.71	39.75	32.02	5.92	42.63
Sum Snappers								0.02				
Sum Groupers								1.76	0.82	0.08		
Sum Grunts								2.16		6.84		
Sum Croakers								2.81	10.61	0.35		
Sum Seabreams								0.31	0.21	0.24	8.80	
Sum Sharks								0.28	0.03	0.52	2.44	
Sum Rays								0.72	0.10	1.00	0.15	
Sum Squids										0.17	0.43	0.42
Sum											0.05	2.20
0.02												

Number of stations included in analysis, total and by depth strata

17 3 4 6 4

SWEPT AREA ANALYSIS FROM STATION 987 TO STATION 1088

Guinea 2006

ONLY STATIONS IN SECTOR
ARE INCLUDED

SPECIES NAME	SAMPLE DISTRIB. BY CATCH CLASSES					% inci-dence	Mean dens. t/nm ²	Mean densities by bottom depth strata t/nm ²				
	Lower limits, Kg/nm							- 30m	30- 50m	50-100m	100-400m	
	>0	10	30	100	300	1000						
Trachurus trecae	3	2			1		38	25.93	0.02	0.54	136.30	0.66
Antigonia capros	1	1		1			19	3.69			0.04	14.72
Decapterus punctatus	3	2	1	2			50	3.02	0.52	2.87	9.87	
Saurida brasiliensis	1				1		13	2.18			11.63	
Sardinella aurita	4	2		1			44	2.15	0.07	5.59		0.23
Brachydeuterus auritus	3	1	1	1			31	2.00	2.93	3.87		
Chloroscombrus chrysurus	4		1	1			31	1.85	1.34	4.26	0.01	
Sardinella maderensis	3		1	1			31	1.70	8.03	0.52		
Decapterus rhonchus	7	1	3				50	1.58	2.11	3.17		
Chlorophthalmus atlanticus	1	1		1			19	1.03				4.12
Pseudupeneus prayensis	5	2	1				44	0.75	0.10	1.88	0.15	
Ariommá bondi	2		2				25	0.61		0.03	1.13	1.55
Illlex coindetii	6	1	1				50	0.49			0.02	1.95
Pagrus caeruleostictus	4	3					38	0.39	0.67	0.71		
Pagellus bellottii	6	1	1				50	0.37	0.12	0.93		
Rachycentron canadum	1		1				13	0.20	1.07			
Decapterus & Trachurus JUVENILES			1				6	0.20	1.07			
Acanthurus monroviae				1			6	0.20			0.54	
Lethrinus atlanticus			1					0.16		0.42		
Sphyraena afra	2	1					19	0.13	0.07	0.31		
Aulopus cadenati	1	1					13	0.13				0.51
Merluccius polli	1	1					6	0.11				0.44
Lutjanus goreensis			1					0.09		0.24		
Synagrops microlepis	2	1					19	0.08		0.01		0.30
Aluterus heudelotii	2						13	0.07		0.17		
Engraulis encrasicolus	2						13	0.07	0.40			
Sardinella spp. (juv.)		1					6	0.07	0.37			
Sepia officinalis hierredda	8						44	0.06	0.14	0.08		
MYCTOPHIDAE	2						13	0.05		0.10		0.06
Bodianus speciosus	1							0.05		0.14		
Parapenaeus longirostris	2						6	0.01				0.03
Nematopalaemon hastatus	1						6		0.34	0.71	0.43	1.23
Shrimps, small, non comm.	1							0.67				
Other fish												
Sum all species							50.09	19.37	27.09	159.81	25.57	
Sum Snappers							0.16			0.44		
Sum Groupers							0.03			0.07		
Sum Grunts							2.01	2.93		3.88		
Sum Croakers												
Sum Seabreams							0.77	0.79		1.66		
Sum Sharks							0.08					
Sum Rays							0.04		0.07	0.02		0.07
Sum Squids							0.56	0.14	0.08	0.02		1.99
Sum												

Number of stations included in analysis, total and by depth strata

16 3 6 3 4

SWEPT AREA ANALYSIS FROM STATION 987 TO STATION 1063

Sierra Leone 2006

ONLY STATIONS IN SECTOR
ARE INCLUDED

SPECIES NAME	SAMPLE DISTRIB. BY CATCH CLASSES					% inci-dence	Mean dens. t/nm ²	Mean densities by bottom depth strata t/nm ²				
	Lower limits, Kg/nm							- 30m	30- 50m	50-100m	100-200m	
	>0	10	30	100	300	1000						
Ariomma bondi	2	1			1		16	7.87		24.08	4.04	
Chloroscombrus chrysurus	7	1	3	2			52	6.96	12.92	7.80	1.02	
Priacanthus arenatus	9	1	2	2			56	6.18	0.47	0.25	18.58	
Decapterus punctatus	6	2	2	1	1		44	3.59	0.05	1.96	9.23	
Pagellus bellottii	7	2	4				52	1.27	1.67	1.29	0.98	
Pagrus caeruleostictus	11	6		1			68	1.21	2.28	0.95	0.54	
Brachydeuterus auritus	7	3	3				52	0.96	1.01	1.87	0.11	
Sardinella aurita	4	3	3				40	0.83		0.78	1.80	
Dentex congensis	2	1	1	1			20	0.81			2.48	
Decapterus rhonchus	11	1		1			48	0.76	0.37	0.02	1.98	
Sphyraena guachancho	9	3	1				36	0.51	0.26	1.34		
Ilisha africana	3	1	2				24	0.42	0.69	0.64		
Galeoides decadactylus	10	2	1				32	0.40	1.03	0.23		
Pseudupeneus prayensis	15	3					72	0.36	0.57	0.36	0.20	
Rhizoprionodon acutus	3		1				16	0.29	0.03	0.88		
Pomadasys rogeri	4		1				16	0.29	0.06	0.85		
Dactylopterus volitans	6	1	1				32	0.29	0.26	0.62	0.05	
Balistes capriscus	5		2				28	0.28	0.45	0.42	0.01	
Selene dorsalis	11	1					44	0.27	0.34	0.49	0.01	
Pteroscion peli	4	1	1				24	0.25	0.29	0.50		
Pseudotolithus senegalensis	11	1					28	0.21	0.48	0.18		
Trachurus trecae	4	2					24	0.21		0.01		
Sardinella maderensis	10	1					36	0.20	0.07	0.18	0.36	
Engraulis encrasicolus	1		1				8	0.19		0.60		
Trichiurus lepturus	5	1					24	0.14	0.19	0.26		
Pentanemus quinquarius	3	1					16	0.11	0.32	0.02		
Lagocephalus laevigatus	9						36	0.08	0.04		0.18	
Dentex angolensis	4	1					20	0.08		0.06	0.26	
Chromis cadenati	1	1					8	0.08			0.25	
Parapenaeopsis atlantica	1	1					8	0.07	0.21			
Epinephelus aeneus	4						16	0.06		0.05		
Lethrinus atlanticus	5	1					24	0.06	0.02	0.17		
Caranx cryos	10						36	0.06	0.14	0.06		
Alectis alexandrinus	3	1					12	0.06		0.18		
Acanthurus monroviae	3						12	0.06		0.18		
Boops boops	4						16	0.05				
Scomber japonicus	2						8	0.05				
Chaetodipterus goreensis	5						20	0.05	0.07	0.10		
Balistes punctatus	3						12	0.05		0.14		
Penaeus notialis	6						24	0.03	0.03	0.05		
Penaeus kerathurus	4						16	0.01	0.01			
Shrimps, small, non comm.	1						4	0.79	1.26	0.75	0.52	
Other fish											0.82	
Sum all species								36.50	25.59	24.18	63.70	5.52
Sum Snappers								0.03	0.09	0.01	0.01	
Sum Groupers								0.06		0.06	0.14	
Sum Grunts								1.29	1.19	2.72	0.11	
Sum Croakers								0.49	0.83	0.68	0.03	
Sum Seabreams								3.43	3.95	2.25	4.44	0.38
Sum Sharks								0.32	0.04	0.88	0.08	
Sum Rays								0.05	0.05	0.03	0.06	0.04
Sum Squids								0.04	0.01	0.09	0.02	0.01
Sum								0.05				

Number of stations included in analysis, total and by depth strata

25 8 8 8 1

SWEPT AREA ANALYSIS FROM STATION 987 TO STATION 1088

Liberia 2006

ONLY STATIONS IN SECTOR
ARE INCLUDED

SPECIES NAME	SAMPLE DISTRIB. BY CATCH CLASSES					% inci-dence	Mean dens. t/nm ²	Mean densities by bottom depth strata t/nm ²			
	Lower limits, Kg/nm							- 30m	30- 50m	50-100m	100-200m
	>0	10	30	100	300	1000					
Pseudotolithus typus		1	1				1.38		5.81		
Trachurus trecae	1	1	3				1.09			1.64	
Dentex congensis	3	2	3				3.8	1.03		1.54	
Priacanthus arenatus	11	1		1			6.2	1.01	0.01	1.52	
Sphyraena guachancho	17	2	2				6.2	0.92	0.43	2.80	0.31
Dentex angolensis	7	8					6.7	0.80			1.21
Boops boops	6	1	2				4.3	0.58			0.87
Pseudotolithus senegalensis	6	5					3.3	0.38	1.11	1.15	
Brachydeuterus auritus	8	3					5.2	0.34	0.27	0.62	0.25
Sardinella aurita	5		1				2.9	0.31			0.46
Ilisha africana	5		1				2.9	0.27	0.12	1.08	
Selene dorsalis	12	1					5.7	0.24	0.19	0.59	0.12
Ariomma bondi	11		1				5.7	0.23			0.34
Squatina oculata	3	2					1.9	0.22			0.33
Pagellus bellottii	11	1					5.2	0.21			0.32
Pteroscion peli	4	1					2.4	0.21	0.45	0.69	
Galeoides decadactylus	8	1					3.8	0.19	0.15	0.73	
Dectapterus punctatus	6	1					2.9	0.14			0.21
Trichiurus lepturus	8	1					4.3	0.12	0.33	0.33	0.02
Sardinella maderensis	7	1					3.8	0.11	0.18	0.39	
Chloroscombrus chrysurus	5	1					2.9	0.11	0.54	0.25	
Dentex canariensis	5						2.4	0.10			0.16
Parapenaeopsis atlantica	4	1					2.4	0.10	0.49	0.24	
Epinephelus aeneus	6						2.9	0.09			0.14
Pomadasys jubelini	5	1					2.9	0.08	0.05	0.30	
Penaeus notialis	7						2.9	0.07	0.03	0.27	
Umbrina canariensis	7						3.3	0.07		0.10	0.07
Lutjanus dentatus		1					5	0.07		0.30	
Arius heudelotii		1					5	0.07		0.31	
Illex coindetii	16						6.7	0.06		0.02	0.08
Scorpaena scrofa	6						2.9	0.06			0.08
Raja miraletus	8						3.8	0.05		0.10	0.03
Parapenaeus longirostris	5						1.9	0.01			0.01
Penaeus kerathurus	2						10		0.02	0.87	0.71
Other fish								0.72	0.86		
Sum all species							11.44	5.22	16.96	10.42	
Sum Snappers							0.07		0.30		
Sum Groupers							0.09			0.14	
Sum Grunts							0.42		0.95		
Sum Croakers							2.10	1.62	7.86	0.25	
Sum Seabreams							2.75			0.12	
Sum Sharks							0.23		0.02	4.15	
Sum Rays							0.07		0.15	0.33	
Sum Squids							0.09		0.03	0.05	
Sum	0.51									0.12	

Number of stations included in analysis, total and by depth strata

21

2

5

14

Annex VI Excel sheet used for calculations of biomass and confidence intervals

This example is the biomass of seabreams in Benin 2002

This sheet is used to calculate stratified mean density, total biomass, and 95% confidence limits on the total biomass.

Inputs are only required in the yellow fields and optionally the t-value can be set. NOTE that the Station field MUST be 1 even if there is no catch

Density (t/nm^2) is from NAN-SIS and Coefficient of variation (CV) is from GRAFER using the same depth intervals.

The underlying assumption is that the CV from the catch (kg/hour) is equal for the density (t/nm^2), i.e. that the swept area is constant per hour.

Equation numbers (1) and (2) refers to Appendix in report

Input from NANSIS					GRAFER				
Depth (m)	Area	No Stations	Density (t/nm^2)	CV (kg/hour)	Equation(1)=		SD	Est. Variance	Equation (2)=
20-30	387	6	0.08	1.83	0.04		0.146	0.021	0.001
31-50	134	6	0.53	1.54			0.816	0.666	0.003
51-100	244	5	2.59	1.20			3.108	9.660	0.197
Total	5561					Var(strat-mean)=		0.20	

$$t\text{-} value = \boxed{2}$$

Stratified mean = 0.96

SE(strat-mean)= 0.45

95% Confidence limits:

Annex VII Instruments and fishing gear used

Echo sounder

The SIMRAD EK500/38 kHz scientific sounder was used during the survey for fish abundance estimation. The lowering keel was not submerged during the survey. The Bergen Echo Integrator system (BEI) was used to scrutinise the acoustic records. System calibration experiment using a standard copper sphere was performed 11.01.2006 The settings of 38 kHz echo sounder were as follows:

Transceiver-1 menu (38 kHz lowering keel)

Transducer depth	5.50 m
Absorbtion coeff.	10 dB/km
Pulse length	medium (1ms)
Bandwidth	wide
Max power	2000 Watt
2-way beam angle	-21.0 dB
SV transducer gain	26.96 dB
TS transducer gain	27.07 dB
Angle sensitivity	21.9
3 dB beamwidth	6.9 dg along / athwardship: 6.8 dg
Alongship offset	-0.07 "
Athwardship offset	0.08 "

Display menu

Echogram	1 (38 kHz)
Sv colour min	-67 dB

Printer- menu

Echogram	1 (38 kHz)
Range	50, 100, 250, 500, 750 and 1000 m
Range start	0
Bottom range	15 m
Bottom range start	10 m
Sv colour min	-67 dB
TVG	20 log R

Bottom detection menu Minimum level -40 dB

Fishing gear

The vessel has "Harstad" and "Åkrahamn" pelagic trawls and "Gisund super bottom trawl".

The bottom trawl has a headline of 31 m, footrope 47 m and 20 mm meshsize in the codend with an innernet of 10 mm meshsize. The estimated opening is 6 m (observed 5.7) and distance between wings during towing about 18 m. The sweeps are 40 m long. The trawl is equipped with a 12" rubber bobbins gear. The doors are of 'Thyborøn' combi type, 7.81 m², 1670 kg, their distance while trawling about 45 - 55 m in average, depending on the depth (least distance at low depths). This distance can be kept constant (about 50 m) at all depths by the use of a 9.5 m strap between the wires at 130 m distance from the doors, normally applied at depths greater than 80 m. On the present survey, however, the strap was not applied because most of the trawl hauls were made in shallower waters.

The SCANMAR system was used on all trawl hauls. This equipment consists of sensors, a hydrophone, a receiver, a display unit and a battery charger. Communication between sensors and ship is based on acoustic transmission. The doors are fitted with sensors to provide information on their distance and a height sensor is fitted to the bottom trawl to measure the trawl opening and provide information on clearance and bottom contact.

The pelagic trawl can be equipped with a trawleye that provides information on the trawl opening and the distance of the footrope to the bottom.

Annex VIII Swept Area Estimates of Main Demersal Groups

Total swept area estimates 2006

Country	Seabreams	Grunts ¹	Croakers	Groupers	Snappers	<i>B. auritus</i>	Sharks	Rays	Cephalopods
Guinea Bissau	15907	142	11736	128	0	11622	1809	1865	3837
Guinea	7873	41	0	287	1804	19823	189	358	1932
Sierra Leone	21362	2131	2897	440	191	5940	2144	298	260
Liberia	13031	369	9535	440	297	1628	1056	306	407
Total	58173	2683	24168	1295	2292	39013	5198	2826	6435

¹ Grunts excluding *Brachydeuterus auritus*

Guinea Bissau, swept area estimates 2006

Depth	20-30 m	30-50 m	50-100 m	100-200 m
Area NM ²	1000	1600	1560	1270

Densities				
Seabreams	10.61	1.14	0.24	2.44
Grunts	0.03	0.07		
Croakers		0.35		8.80
Groupers		0.08		
Snappers				
<i>B. auritus</i>	0.79	6.77		
Sharks	0.21	0.52	0.15	0.42
Rays	0.03	1.00	0.11	0.05
Cephalopods	0.10	0.17	0.43	2.20

	Biomass (tonnes)				Total
Seabreams	10610	1824	374	3099	15907
Grunts	30	112	0	0	142
Croakers	0	560	0	11176	11736
Groupers	0	128	0	0	128
Snappers	0	0	0	0	0
<i>B. auritus</i>	790	10832	0	0	11622
Sharks	210	832	234	533	1809
Rays	30	1600	172	64	1865
Cephalopods	100	272	671	2794	3837
Total	11770	16160	1451	17666	47047

Guinea Bissau, swept area estimates 2006

Depth	20-30 m	30-50 m	50-100 m	100-200 m
Area NM ²	1000	1600	1560	1270

	Densities			
Seabreams	10.61	1.14	0.24	2.44
Grunts	0.03	0.07		
Croakers		0.35		8.80
Groupers		0.08		
Snappers				
<i>B. auritus</i>	0.79	6.77		
Sharks	0.21	0.52	0.15	0.42
Rays	0.03	1.00	0.11	0.05
Cephalopods	0.10	0.17	0.43	2.20

	Biomass (tonnes)				Total
Seabreams	10610	1824	374	3099	15907
Grunts	30	112	0	0	142
Croakers	0	560	0	11176	11736
Groupers	0	128	0	0	128
Snappers	0	0	0	0	0
<i>B. auritus</i>	790	10832	0	0	11622
Sharks	210	832	234	533	1809
Rays	30	1600	172	64	1865
Cephalopods	100	272	671	2794	3837
Total	11770	16160	1451	17666	47047

Sierra Leone, swept area estimates 2006

Depth	20-30 m	30-50 m	50-100 m	100-200 m
Area NM ²	1640	2160	2220	440

	Densities			
Seabreams	3.95	2.25	4.44	0.38
Grunts	0.18	0.85		
Croakers	0.83	0.68	0.03	
Groupers		0.06	0.14	
Snappers	0.09	0.01	0.01	
<i>B. auritus</i>	1.01	1.87	0.11	
Sharks	0.04	0.88	0.08	
Rays	0.05	0.03	0.06	0.04
Cephalopods	0.01	0.09	0.02	0.01

	Biomass (tonnes)				Total
Seabreams	6478	4860	9857	167	21362
Grunts	295	1836	0	0	2131
Croakers	1361	1469	67	0	2897
Groupers	0	130	311	0	440
Snappers	148	22	22	0	191
<i>B. auritus</i>	1656	4039	244	0	5940
Sharks	66	1901	178	0	2144
Rays	82	65	133	18	298
Cephalopods	16	194	44	4	260
Total	10102	14515	10856	189	35663

Liberia, swept area estimates 2006

Depth	0-30 m	30-50 m	50-100 m	100-200 m
Area NM ²	850	990	3140	500

	Densities		
Seabreams		4.15	
Grunts	0.05	0.33	
Croakers	1.62	7.86	0.12
Groupers		0.14	
Snappers		0.30	
<i>B. auritus</i>	0.27	0.62	0.25
Sharks		0.02	0.33
Rays		0.15	0.05
Cephalopods		0.03	0.12

	Biomass (tonnes)			Total
Seabreams	0	0	13031	0
Grunts	43	327	0	0
Croakers	1377	7781	377	0
Groupers	0	0	440	0
Snappers	0	297	0	0
<i>B. auritus</i>	230	614	785	0
Sharks	0	20	1036	0
Rays	0	149	157	0
Cephalopods	0	30	377	0
Total	1649	9217	16202	0
				27068

Annex IX Regional estimates, April-May 2006

April-May 2006: Round sardinella (*Sardinella aurita*), number in millions

Length cm	Guinea Bissau	Guinea	Sierra Leone	Liberia	TOTAL
5					
6					
7					
8		71	3		74
9		915	12	10	937
10		2 674	85	122	2 881
11	603	3 535	247	48	4 434
12	1 231	4 266	399	27	5 923
13	327	2 783	464	22	3 596
14	50	1 052	490	45	1 637
15		1 668	192	71	1 932
16	123	1 587	41	85	1 836
17	246	2 838	24	134	3 242
18	921	970	8	34	1 932
19	860	272	29	6	1 167
20	61	299	0	4	365
21		245	101	2	349
22		27	7	2	36
23		320			320
24					
25					
26					
27					
28					
29					
30					
31					
32					
33					
34					
35					
36					
37					
38					
39					
40					
41					
42					
43					
44					
45					
46					
47					
48					
49					
50					
Total	4 421	23 523	2 105	612	30 661

Annex IX continued

April-May 2006: Round sardinella (*Sardinella aurita*), biomass in tonnes

Length cm	Guinea Bissau	Guinea	Sierra Leone	Liberia	TOTAL
5					
6					
7			19		19
8		409	94	81	584
9		7 376	928	1 327	9 631
10		29 095	3 538	691	33 323
11	8 618	50 536	7 332	501	66 988
12	22 597	78 316	10 740	519	112 172
13	7 552	64 357	14 054	1 277	87 240
14	1 440	30 148	6 732	2 499	40 818
15		58 393	1 720	3 598	63 711
16	5 185	67 033	1 213	6 767	80 198
17	12 373	142 960	449	2 002	157 784
18	54 817	57 749	2 036	393	114 995
19	59 915	18 975	14	291	79 196
20	4 972	24 251	9 479	191	38 894
21		22 890	788	173	23 851
22		2 915			2 915
23		39 076			39 076
24					
25					
26					
27					
28					
29					
30					
31					
32					
33					
34					
35					
36					
37					
38					
39					
40					
41					
42					
43					
44					
45					
46					
47					
48					
49					
50					
Total	177 470	694 478	59 137	20 310	951 395

Annex IX continued

April-May 2006: Flat sardinella (*Sardinella maderensis*), numbers in millions

Length cm	Guinea Bissau	Guinea	Sierra Leone	Liberia	TOTAL
5				1	1
6			10		10
7		66	38	5	110
8		1 177	104	7	1 289
9		2 604	51	21	2 676
10		482	21	20	522
11		212	44	12	269
12		242	69	6	318
13		273	31	4	307
14		718	19	3	740
15		556	21	2	579
16		1 044	44	0	1 088
17		1 427	93	1	1 521
18		373	62		435
19		1 016	74	1	1 090
20		646	73		720
21		177	41		218
22			20		20
23		54	7	3	64
24		3	14	8	25
25			1	3	4
26			1	5	6
27					
28				1	1
29					
30					
31					
32					
33					
34					
35					
36					
37					
38					
39					
40					
41					
42					
43					
44					
45					
46					
47					
48					
49					
50					
Total		11 071	837	103	12 012

Annex IX continued

April-May 2006: Flat sardinella (*Sardinella maderensis*), biomass in tonnes

Length cm	Guinea Bissau	Guinea	Sierra Leone	Liberia	TOTAL
5				1	1
6			27		27
7		271	157	21	449
8		7 013	622	42	7 676
9		21 655	427	172	22 254
10		5 412	231	220	5 862
11		3 131	652	182	3 965
12		4 594	1 316	118	6 028
13		6 509	731	95	7 335
14		21 232	570	93	21 895
15		20 078	766	67	20 911
16		45 481	1 900	13	47 394
17		74 190	4 816	40	79 047
18		22 904	3 816		26 720
19		73 090	5 291	41	78 422
20		54 012	6 125		60 137
21		17 074	3 942		21 015
22			2 167		2 167
23		6 775	864	396	8 035
24		462	1 944	1 197	3 603
25			179	506	685
26			115	947	1 061
27					
28				236	236
29					
30					
31					
32					
33					
34					
35					
36					
37					
38					
39					
40					
41					
42					
43					
44					
45					
46					
47					
48					
49					
50					
Total		383 883	36 656	4 388	424 927

Annex IX continued

April-May 2006: Anchovy (*Engraulis encrasicolus*), numbers in millions

Length cm	Guinea Bissau	Guinea	Sierra Leone	Liberia	TOTAL
5		0	160	42	202
6		0	169	44	214
7		132	152	39	323
8		362	71	19	452
9		526	9	2	537
10		132	9	2	143
11		33			33
12		0			0
13					0
14					0
15					0
16					0
17					0
18					0
19					0
20					0
Total		1 184	571	149	1 903

April-May 2006: Anchovy (*Engraulis encrasicolus*), biomass in tonnes

Length cm	Guinea Bissau	Guinea	Sierra Leone	Liberia	TOTAL
5			144	38	182
6			251	65	317
7		300	345	90	735
8		1 200	237	62	1 498
9		2 436	41	11	2 488
10		822	56	15	893
11		270			270
12					
13					
14					
15					
16					
17					
18					
19					
20					
Total		5 028	1 074	280	6 382

Annex IX continued

April-May 2006: Ilisha (P1), number in millions

Length cm	Guinea Bissau	Guinea	Sierra Leone	Liberia	TOTAL
5				1	1
6	122		2		124
7	951		20	2	973
8	922		22	6	949
9	1 331		28	2	1 361
10	833		20	4	857
11	151		4	0	156
12	218		5	2	225
13	362		11		372
14	933		25	5	963
15	730		22	2	754
16	796		22	7	826
17	453		20	13	487
18	516		17	1	534
19	449		16	2	468
20	144		8		151
21	81		4	2	87
22	122		2	0	124
23					
24					
25					
26					
27					
28					
29					
30					
Total	9 113		248	49	9 411

Annex IX continued

April-May 2006: Ilisha (P1), biomass in tonnes

Length cm	Guinea Bissau	Guinea	Sierra Leone	Liberia	TOTAL
5				1	1
6	267		5		272
7	3 209		68	7	3 284
8	4 529		109	27	4 665
9	9 127		191	16	9 334
10	7 716		182	41	7 938
11	1 842		46	6	1 893
12	3 405		80	25	3 510
13	7 117		210		7 327
14	22 764		609	112	23 484
15	21 746		659	68	22 473
16	28 623		796	247	29 666
17	19 432		873	558	20 863
18	26 132		865	58	27 055
19	26 657		971	122	27 750
20	9 903		518		10 421
21	6 441		288	163	6 891
22	11 073		226	21	11 320
23					
24					
25					
26					
27					
28					
29					
30					
Total	209 981		6 695	1 471	218 147

Annex IX continued

April-May 2006: Cunene horse mackerel (*Trachurus trecae*), numbers in millions

Length cm	Guinea Bissau	Guinea	Sierra Leone	Liberia	TOTAL
5	0				0
6	3				3
7	13				13
8	11	12			23
9		148			148
10		292	1		293
11		48	4		52
12		17	2	7	26
13		9	1	22	32
14	1			6	6
15				6	6
16	1			31	32
17	9			27	36
18	19			9	28
19	18			2	20
20	4				4
21	2				2
22	1				1
23	1				1
24	1				1
25					
26					
27					
28					
29					
30		10	5		15
31		7	3		10
32					
33					
34					
35					
36					
37					
38					
39					
40					
41	6				6
42	9				9
43	10				10
44	10				10
45	11				11
46	8				8
47	1				1
48	1				1
49					
50	1				1
Total	139	542	17	110	808

Annex IX Regional estimates, April-May 2006

April-May 2006: Round sardinella (*Sardinella aurita*), number in millions

Length cm	Guinea Bissau	Guinea	Sierra Leone	Liberia	TOTAL
5					
6					
7					
8		71	3		74
9		915	12	10	937
10		2 674	85	122	2 881
11	603	3 535	247	48	4 434
12	1 231	4 266	399	27	5 923
13	327	2 783	464	22	3 596
14	50	1 052	490	45	1 637
15		1 668	192	71	1 932
16	123	1 587	41	85	1 836
17	246	2 838	24	134	3 242
18	921	970	8	34	1 932
19	860	272	29	6	1 167
20	61	299	0	4	365
21		245	101	2	349
22		27	7	2	36
23		320			320
24					
25					
26					
27					
28					
29					
30					
31					
32					
33					
34					
35					
36					
37					
38					
39					
40					
41					
42					
43					
44					
45					
46					
47					
48					
49					
50					
Total	4 421	23 523	2 105	612	30 661